

# The Acupuncture Evidence Project

A Comparative Literature Review

John McDonald

Stephen Janz

January 2017

(Revised Edition)

Commissioned by  
Australian Acupuncture and Chinese Medicine Association Ltd



## **Acknowledgments**

The Acupuncture Evidence Project: A Comparative Literature Review was funded by the Australian Acupuncture and Chinese Medicine Association Ltd (AACMA).

Research and analysis was undertaken by John McDonald PhD. The Project was initiated and managed by Stephen Janz.

Tables 3, 4, 5, and 6, and Sections 2 and 3 were written by John McDonald. Stephen Janz wrote the preface and the authors collaborated on Section 1, Tables 7 and 8, and the Appendix. Research design by Stephen Janz and John McDonald.

The authors thank Professor Caroline Smith, National Institute of Complementary Medicine, Western Sydney University, for providing advice regarding evidence levels for assisted reproduction trials; Associate Professor Zhen Zheng, RMIT University, for identifying the evidence levels for post-operative nausea and vomiting and post-operative pain; Judy James for technical and editorial assistance; and the AACMA board for funding and supporting this project. The authors also thank Dr Suzanne Cochrane, Western Sydney University; Associate Professor Chris Zaslowski, University of Technology Sydney; and Associate Professor Zhen Zheng, RMIT University, for providing pre-publication commentary and advice.

## **Conflicts of interest**

Dr John McDonald was a co-author of three of the research papers referenced in this review. Professor Caroline Smith was a co-author of six of the research papers referenced in this review, and Associate Professor Zhen Zheng was co-author of one of the research papers in this review. There were no other conflicts of interest.

The Acupuncture Evidence Project is a publication of the Australian Acupuncture and Chinese Medicine Association Ltd.

© Australian Acupuncture and Chinese Medicine Association Ltd 2016, 2017. First published December 2016. Revised and updated 17 January 2017.

This document should be cited as: McDonald J, Janz S. The Acupuncture Evidence Project: A Comparative Literature Review (Revised edition). Brisbane: Australian Acupuncture and Chinese Medicine Association Ltd; 2017. <http://www.acupuncture.org.au>.

## **Electronic documents**

This work is copyright. You may download, display, print and reproduce this material in unaltered form only (retaining this notice) for your personal, non-commercial use, or use within your organisation. Apart from any use as permitted under the Copyright Act 1968, all other rights are reserved.

ISBN 978-0-9954289-3-5

Requests for further authorisation should be directed to: The Australian Acupuncture & Chinese Medicine Association Ltd. PO Box 1635 COORPAROO DC QLD 4151 or via <http://www.acupuncture.org.au>.

## **Paper-based publication**

This work is copyright. Apart from any use permitted under the Copyright Act 1968, no part may be reproduced by any process without written permission from the Australian Acupuncture and Chinese Medicine Association Ltd.

ISBN 978-0-9954289-2-8

Requests and inquiries concerning reproduction and rights should be addressed to: Australian Acupuncture and Chinese Medicine Association Ltd. PO Box 1635 COORPAROO DC QLD 4151 or via <http://www.acupuncture.org.au>.

## **Disclaimer**

This document aims to review the best available evidence to assist in decision-making. The authors and the Australian Acupuncture and Chinese Medicine Association Ltd give no warranty that the information contained in this document and any online updates available on the AACMA website is correct or complete.

Neither the authors nor the AACMA shall be liable for any loss whatsoever whether due to negligence or otherwise arising from the use of or reliance on this document. This document can be downloaded from the AACMA website. <http://www.acupuncture.org.au>.

## **ABSTRACT**

### **Background**

The acupuncture evidence project investigated the state of the evidence regarding acupuncture, with the focus on systematic reviews and meta-analyses. The Australian Department of Veterans' Affairs 2010 Alternative Therapies Review and United States Department of Veterans Affairs Acupuncture Evidence Map 2014 were used as baselines, then evidence levels were updated to reflect subsequent research.

### **Methods**

A search of PubMed and Cochrane Library for systematic reviews and meta-analyses from March 2013 to September 2016 was conducted. Three reviews from October 2016 to January 2017 were also included. Evidence levels were graded using NHMRC levels. Risk of bias was assessed using the Cochrane GRADE system where possible. All results were displayed in tables to demonstrate changes in evidence level over time, as well as the current state of evidence by clinical area.

### **Results**

Of the 122 conditions reviewed, evidence of effect was found at various levels for 117 conditions. Five conditions were assessed as 'no evidence of effect'. The level of evidence had increased for 24 conditions since the previous reviews. The findings of this review are limited by the mounting evidence that sham/placebo controls used in acupuncture trials are not inert, which is likely to lead to a consistent underestimation of the true effect size of acupuncture interventions.

### **Conclusions**

Systematic reviews published up to January 2017 indicate that acupuncture has a 'positive effect' on eight conditions (migraine prophylaxis, headache, chronic low back pain, allergic rhinitis, knee osteoarthritis, chemotherapy-induced nausea and vomiting, post-operative nausea and vomiting and post-operative pain), 'potential positive effect' on a further 38 conditions, 'unclear/insufficient evidence' for 71 conditions and 'no evidence of effect' for five conditions. Evidence of cost-effectiveness was identified for 10 conditions, and evidence for safety was identified for nine conditions.

## **ADDENDUM TO REVISED EDITION**

The Acupuncture Evidence Project: A Comparative Literature Review was released in limited circulation in December 2016. New research is regularly being reported and there is always a risk that a significant paper might be published just after a review is completed. Just prior to general publication, the authors became aware of three new studies with findings that were relevant to the review's results. As the objective of the review was to identify the state of acupuncture research, the authors decided to incorporate these late papers and amend the results accordingly, even though the new papers fell outside the initial search dates. These papers concerned chronic pain and assisted reproduction.

The authors have decided to leave the table headers and narrative to reflect evidence levels to 2016. This is a more accurate reflection of the study's results, as only one study has been included from 2017. The additional studies are clearly indicated in the tables. A plain English summary has been added to the Appendix in this revised edition.

17 January 2017

## **PREFACE**

### **Background**

Evidence of effectiveness underpins the validity of all health care interventions. Acupuncture has been practised for thousands of years; however, research into its effectiveness and cost effectiveness is in its relative infancy. The first significant attempt to identify the evidence validating the role of acupuncture was undertaken by the World Health Organization (WHO) in 1979 (1). WHO conducted a Delphi-like symposium in Beijing in 1979 where physicians from around the world identified 43 diseases which they believed acupuncture may benefit (1). The 1979 report was criticised because it was not based on clinical trials, rather the clinical experience of the participants (2).

The US Department of Health and Human Services National Institutes of Health (NIH) released a consensus statement on acupuncture in 1997 (3). The statement identified the efficacy of acupuncture for adult postoperative and chemotherapy nausea and vomiting, and for postoperative dental pain. In addition, the statement found support for the use of acupuncture for a range other conditions and identified the need for further research (3).

In 1996, WHO held a consultation on acupuncture in Cervia, Italy. At this meeting it was decided to review acupuncture again, this time focusing on the now-numerous controlled clinical trials. The result was a review of clinical trials up until early 1999 and culminated in 'Acupuncture: review and analysis of controlled clinical trials' published by WHO in 2002 (4). The report identified 28 conditions for which acupuncture was found to be effective, and nearly 100 others where there was a therapeutic effect (4). It was acknowledged at the time that there were problems with the methodology in some of the trials included in the review (4). Notwithstanding these limitations, the WHO report indicated a growing and convincing body of evidence indicating that there was more to acupuncture than the placebo effect (4).

### **The Acupuncture Evidence Project**

Fourteen years after the WHO publication on acupuncture evidence there has been further refinement in the conduct of clinical trials, not just for acupuncture, but in health-care generally, with clearer guidelines on how to rate the quality of the evidence. The Australian Acupuncture and Chinese Medicine Association Ltd (AACMA) identified the need for an updated review of the literature with greater rigour than was possible in the past, and commissioned the acupuncture evidence project. AACMA engaged experienced clinician and researcher, Dr John McDonald PhD, to conduct the research and analysis for the project.

This review draws on two prior comprehensive literature reviews, one conducted for the Australian Department of Veterans' Affairs (DVA) in 2010 and another conducted for the United States Department of Veterans Affairs (USVA) in 2013 (5, 6). The research identified by these reviews was pooled, then a search of further literature from 2013 to 2016 was conducted. Trials were assessed using the National Health and Medical Research Council (NHMRC) levels of evidence, with risk of bias assessed using the Cochrane GRADE system (7, 8). Results have been tabulated to indicate not just the current state of the evidence, but to indicate how the quality and quantity of evidence has changed from 2005 to 2016. In this review, 122 conditions across 14 broad clinical areas were identified and, of these, only five conditions found 'no evidence of effect' for acupuncture. The level of evidence was found by this review to have increased for 24 conditions.

### **How to use this review**

This project sets a new benchmark to inform acupuncturists, the public, researchers, health departments, governments, and other health providers that acupuncture has a valuable contribution to make to global healthcare and to assist in reducing the global burden of disease.

Acupuncturists should also take confidence from this report that their clinical expertise has been validated, and to confidently offer their services alongside other health professionals. Students and clinicians can use this report to identify areas of clinical interest which they may have overlooked. Researchers can find inspiration for areas of future investigation where the evidence is currently unclear. This review should also encourage educational institutions to maintain robust programs of study in acupuncture to continue to produce graduates capable of the broad scope of practice that this report indicates. It is no longer possible to say that the effectiveness of acupuncture can be attributed to the placebo effect or that it is useful only for musculoskeletal pain.

*It is no longer possible to say that the effectiveness of acupuncture can be attributed to the placebo effect or that it is useful only for musculoskeletal pain.*

The realisation that pharmacological and surgical interventions are not without their limitations has increased interest in drug-free treatments such as acupuncture (9-13). This review found eight conditions where acupuncture may be used to reduce reliance on pharmacological or surgical options. **Migraine** and **tension headaches** lead to loss of productivity and quality of life; a drug-free therapy has a major health impact and potential cost savings as well as maintaining participation in the workforce (14). Medication is not always an effective or acceptable therapy for **allergic rhinitis** and acupuncture improves the range of interventions available to improve quality of life (15). **Post-operative nausea and vomiting** and **post-operative pain** complicate post-operative management, with acupuncture offering another avenue to enhance care in the post-operative period and reduce reliance on medication alone (16, 17).

**Knee osteoarthritis** is on the increase globally and contributes not just to disability adjusted life years (DALYs), but is an increasing burden on health budgets (18, 19). The option of a safe, drug-free treatment that may improve quality of life and potentially delay surgical intervention has significant potential to control these spiralling costs and DALYs. **Low back pain** is a WHO priority disease, and is the single largest contributor to disability worldwide (20). The finding that acupuncture benefits chronic low back pain is arguably the most important finding from this report. Finally, **chemotherapy induced nausea and vomiting** is an unwanted complication of cancer treatment and is often not fully controlled even with state-of-the-art antiemetics. Acupuncture can assist in improving quality of life for these cancer patients (21).

It has been estimated that there is a 17-year time lag in translating clinical research into clinical practice (22). During this time patients are being deprived of the benefit of a proven therapy. Health policy makers now have eight clear conditions associated with a significant burden of disease where acupuncture should be integrated into current clinical guidelines without further delay. Placebo controlled clinical trials consistently underestimate the true effect size of acupuncture (as discussed in section 1.4), yet they have still demonstrated NHMRC Level 1 evidence for the effectiveness of acupuncture for a further 109 conditions. This review has found a significant improvement in both the quality of studies and the levels of evidence supporting acupuncture since the most recent reviews conducted by the Australian and US Departments of Veterans Affairs.

### **Stephen Janz MPH**

Project Director

The AACMA Acupuncture Evidence Project

Brisbane, Australia

## TABLE OF CONTENTS

<b>ACKNOWLEDGEMENTS, CONFLICTS OF INTEREST</b>	i
<b>ABSTRACT</b>	ii
<b>ADDENDUM TO REVISED EDITION</b>	ii
<b>PREFACE</b>	iii
<b>GLOSSARY OF TERMS</b>	vii
<b>1 INTRODUCTION</b>	1
1.1 Methods	1
1.2 Assessment of the quality of evidence	2
1.3 Efficacy versus effectiveness	3
1.4 Problems with placebo controls in acupuncture trials	3
1.5 Acupuncture recommendations in clinical practice guidelines	4
1.6 Interpreting the findings	5
1.7 Limitations	6
<b>2 CONDITIONS NOW RATED AS ‘EVIDENCE OF POSITIVE EFFECT’ IN THIS REVIEW:</b>	7
2.1 Migraine prophylaxis	7
2.2 Headache (chronic tension-type and chronic episodic)	7
2.3 Low back pain	7
2.4 Knee osteoarthritis pain	8
2.5 Allergic rhinitis (seasonal and perennial/persistent)	8
2.6 Chemotherapy-induced nausea and vomiting (CINV)	8
2.7 Post-operative nausea and vomiting (PONV)	9
2.8 Post-operative pain	9
<b>3 RESEARCH INTO THE MECHANISMS OF ACUPUNCTURE</b>	10
<b>TABLES</b>	
Table 1. Summary of levels of evidence used in this review	3
Table 2. Statistical summary of findings of this review	5

Table 3. Summary of effectiveness/efficacy in acupuncture research literature sorted by evidence levels	11
Table 4. Summary of changes in evidence levels from 2005 to 2016	14
Table 5. Evidence levels from 2005 to 2016: explanatory notes with references	18
Table 6. All conditions reviewed sorted by clinical areas	34
Cardiovascular/respiratory conditions	34
Musculoskeletal disorders	35
Neurological disorders	38
Mental Health	39
Gynaecology and obstetrics	40
Gastrointestinal disorders	43
Headache and migraine	43
Genitourinary/sexual disorders	44
Surgery	45
Oncology	46
Eye, ear, nose, throat	48
Paediatrics	49
Dermatology	49
Miscellaneous conditions	50
Table 7. Conditions reported in this review with evidence of cost-effectiveness	51
Table 8. Conditions reported in this review with evidence of safety	52
<b>APPENDIX</b>	53
Table 9. NHMRC Evidence Hierarchy: designations of 'levels of evidence' according to type of research question	53
GRADE and guideline development	54
Table 10. Cochrane GRADE: Significance of the four levels of evidence	54
Plain English Summary	55
Bottom Line	55
Background	55
Key Results	55
Summary of Findings	56
<b>REFERENCES</b>	60

## GLOSSARY OF TERMS

AACMA	Australian Acupuncture and Chinese Medicine Association Ltd
ART	Assisted reproduction technology
CGRP	calcitonin gene-related peptide
DALYs	disability adjusted life years
DVA	Department of Veterans' Affairs
EA	electroacupuncture
ECP	eosinophilic cationic protein
FEV1	forced expiratory volume in one second
FEV1/FVC	ratio of FEV1 to forced ventilation capacity
GABA	gamma-amino-butyric-acid
GRADE	Grades of Recommendation, Assessment, Development and Evaluation
IgE	immunoglobulin E
IL	interleukin
MA	meta-analysis
MMSE	Mini Mental State Examination
NAD	neck pain and associated disorders
NHMRC	National Health and Medical Research Council
NIH	National Institutes of Health
NSAIDS	non-steroidal anti-inflammatory drugs
OHNSF	Otolaryngology Head Neck Surgery Foundation
PAR	persistent (perennial) allergic rhinitis
PCOS	Polycystic ovarian syndrome
QoL	quality of life
RCT	randomised controlled trial
RQLQ	rhinoconjunctivitis quality of life questionnaire (Juniper)
SAR	seasonal (intermittent) allergic rhinitis
SP	substance P
SR	systematic review
TEAS	transcutaneous electrical acupoint stimulation
TRPV1	transient receptor potential vanilloid 1
USVA	United States Department of Veterans Affairs
VIP	vasoactive intestinal peptide
WAD	whiplash associated disorders

## 1 INTRODUCTION

Increasing interest in acupuncture has led to the need for an updated review of its efficacy, effectiveness and cost effectiveness. Rather than starting from scratch, this review draws on two prior comprehensive literature reviews into acupuncture. The first was undertaken for the Australian Department of Veterans' Affairs (DVA) in 2010, the second for the United States Department of Veterans Affairs (USVA) in 2013 (5, 6). These reviews did not limit themselves to veteran-specific health issues but considered the evidence regarding acupuncture in the broadest health terms. A search was then conducted for studies after the USVA 2013 review and the evidence analysed, graded and tabulated. The advantage of this comparative approach is not only a pragmatic way of avoiding duplication of work; it also demonstrates the increasing quality and quantity of evidence supporting the effectiveness of acupuncture over the eleven-year period (2005 to 2016) from which studies were drawn.

Table 3 categorises the research literature by evidence levels. This table clearly demonstrates the increase in the volume and scope of research on acupuncture since the prior reviews. Table 4 summarises the changes in evidence levels among the reviews. Of note is the finding that eight conditions are now rated as 'evidence of positive effect'. Two of these had been listed as 'unclear' evidence and three as 'evidence of potential positive effect' prior to this review. Table 5 provides brief notes for each condition and the references justifying the change in evidence levels found by this review. Table 6 organises conditions by clinical areas, allowing the reader to get a snapshot of the state of the evidence for any particular clinical area, and identifies areas where further research needs to be undertaken.

Numerous clinical guidelines now recommend acupuncture for a range of conditions. Of the 122 conditions examined in this review, levels of evidence have increased for 24 conditions. There is now 'evidence of positive effect' for eight conditions and 'evidence of potential positive effect' for a further 38 conditions. Seventy-one conditions are rated as 'unclear/insufficient evidence' and only five conditions are currently rated at 'no evidence of effect'. Further statistics can be found at Table 2. The comparative approach used in this study serves as a reminder that a finding of 'no evidence of effect' or 'unclear' does not mean 'ineffective'. Many of the conditions currently rated as 'unclear' have consistently positive findings in systematic reviews, but because they represent new clinical areas for acupuncture research, the evidence is not yet sufficient to allow firm conclusions to be drawn on effectiveness or efficacy. Section 1.6 provides further assistance in interpreting the findings of this review. There are methodological challenges in investigating acupuncture such as the problems with placebo controlled trials (outlined in section 1.4) which are slowly being overcome.

### 1.1 Methods

A review of alternative therapies was published by the Australian Government Department of Veterans' Affairs in 2010 (5). In 2014, the US Department of Veterans Affairs published an Evidence Map of Acupuncture reviewing acupuncture research published in or before March 2013 (6). This review focuses on new evidence between March 2013 and September 2016. Like the two previous reviews, this review concentrates on systematic reviews (SR) and meta-analyses (MA), using Cochrane Systematic Reviews where available. The primary focus of this review is to examine evidence supporting the effectiveness and efficacy of acupuncture (see section 1.3 Efficacy versus effectiveness). Where available, evidence on cost-effectiveness and safety has been included (see Tables 7 and 8).

A search was undertaken on PubMed and the Cochrane Library using the search term 'acupuncture' with limits set for dates between March 2013 and September 2016, and restricted to reviews. All languages were included. Systematic reviews, meta-analyses, network meta-analyses, overviews of systematic reviews (NHMRC level I evidence) and some narrative reviews were included, but protocols for systematic reviews were excluded. Systematic reviews of non-invasive or nonpharmacological interventions or of complementary and/or alternative medicine (CAM) interventions were included if they included acupuncture studies. One systematic review from the Australian Journal of Acupuncture and Chinese Medicine was included although this journal is not included in PubMed listings. A check in January 2017 identified three more relevant SRs and MAs (one from October 2016, one from December 2016, and one from January 2017) which were included. A total of 136 systematic reviews, including 27 Cochrane systematic reviews were included in this review, along with three network meta-analyses, nine reviews of reviews and 20 other reviews. Meta-analyses were conducted for 62 of the non-Cochrane systematic reviews. This review includes pooled data from more than 1,000 randomised controlled trials. Some of the included systematic reviews included studies which were not randomised controlled trials.

## **1.2 Assessment of the quality of evidence**

The Australian DVA review (2010) used the National Health and Medical Research Council (NHMRC) levels of evidence (I-IV) which define the source of the evidence, with Level I being 'evidence obtained from a systematic review of all relevant randomised controlled trials', and Level II being 'evidence obtained from at least one properly designed randomised controlled trial' (7). By this definition, this review examined mainly Level I evidence as systematic reviews and meta-analyses were prioritised. Level II evidence from individual randomised controlled trials has been included occasionally where new high quality randomised trials may have changed the conclusions from the most recent systematic review.

Risk of bias in randomised controlled trials has mainly been assessed by the included systematic reviews using the Cochrane GRADE system of low, unclear or high risk of bias across a number of domains (8). The quality of evidence has also been assessed principally using the GRADE system with randomised controlled trials being assessed as high, moderate, low or very low quality evidence. See the Appendix, Table 10 for an outline of GRADE levels. Where systematic reviews have used other systems of quality assessment (such as PRISMA or Jadad) an attempt has been made to convert these scores to an equivalent within the GRADE system. Some systematic reviews have not reported an assessment of quality of evidence of included trials, and due to time constraints, this review has not attempted to make such an assessment.

In the US Veterans Affairs evidence map, evidence has been assessed at four levels: evidence of positive effect, evidence of potential positive effect, unclear evidence and evidence of no effect. In this review this terminology has largely been adopted with the exception that 'evidence of no effect' has been replaced with 'no evidence of effect' which seems to be more strictly accurate. 'Unclear evidence' was described as conflicting evidence between reviews or between authors within a review, with reviewers summarising the evidence as inconclusive. 'Evidence of potential positive effect' refers to reviews reporting all individual RCTs or pooled effects across RCTs as positive, however the reviewers deeming the evidence insufficient to draw firm conclusions. 'Evidence of positive effect' refers to reviews with consistent statistically significant positive effects and where authors have recommended the intervention (6).

<b>Table 1. Summary of levels of evidence used in this review</b>		
<b>Level</b>	<b>Description</b>	<b>GRADE level (8)</b>
Evidence of positive effect	Reviews with consistent statistically significant positive effects and where authors have recommended the intervention. Strong positive evidence.	Moderate or high quality
Evidence of potential positive effect	Reviews reporting all individual RCTs or pooled effects across RCTs as positive, but the reviewers deeming the evidence insufficient to draw firm conclusions. Moderate positive evidence.	Moderate or high quality
Unclear/insufficient evidence	Reviews consisted mostly of weak positive evidence or conflicting evidence between reviews or between authors within a review, with reviewers summarising the evidence as inconclusive.	Low or very low quality; or conflicting levels of evidence within or between reviews
No evidence of effect	Reviews have consistently found little support for acupuncture.	Consistently low or very low quality

There is not necessarily an exact equivalence between the ‘insufficient evidence’ category used in the Australian DVA review and ‘unclear’ in the USVA evidence map, as some conditions rated as ‘insufficient evidence’ in the former may appear in the ‘evidence of potential positive effect’ category in the latter. Generally speaking, conditions rated as positive or potential positive in this review are conditions for which evidence levels were either moderate or high quality evidence according to the GRADE definitions, and conditions with low or very low quality evidence were rated as unclear even when all included RCTs reported positive outcomes. Conditions previously rated as ‘potential positive’ have been down-graded to ‘no evidence of effect’ where subsequent Cochrane reviews have found little support for acupuncture.

### **1.3 Efficacy versus effectiveness**

Efficacy refers to the ability of an intervention to achieve its intended effect in ideal conditions, i.e. a clinical trial, usually with a placebo control. Effectiveness refers to the result of an intervention in real world clinical practice.

The most common method used to assess efficacy is the randomised sham/placebo controlled blinded clinical trial. This methodology is derived from pharmaceutical research where an inert tablet (‘a sugar pill’) is compared to a medication. ‘Efficacy’ in this example is the measure of the effects of the medication on one group of test subjects minus the effects from the sugar pill on the other group of test subjects. The validity of this method of measuring efficacy rests heavily on the assumption that the sham protocol comparator intervention is inert. If the comparator intervention is not inert, this creates difficulties in measuring efficacy accurately, as discussed in the next section.

### **1.4 Problems with placebo controls in acupuncture trials**

It has been observed from the over 8,000 randomised controlled trials which appear in the Cochrane Database of Controlled Clinical Trials that, in studies where acupuncture is compared with no treatment, waitlist or usual care, there is a significantly larger treatment effect than when acupuncture is compared with some form of sham, placebo or minimal acupuncture (23-27). The Society for Acupuncture Research has dubbed this phenomenon a paradox in acupuncture

research (25). There is evidence that many, if not all, the forms of sham/placebo acupuncture treatment protocols used in acupuncture trials to date are not inert, and exert physiological, and possibly also placebo effects, making it extremely difficult (if not impossible) to accurately measure how much of the observed non-specific effects may be attributable to placebo and/or nocebo effects, and how much is due to the placebo not being inert (23, 26-29). Hence a comparison of the effect size of acupuncture when compared to that of placebo acupuncture may result in consistent underestimation of the true effect size (23, 28).

Some acupuncture researchers are now suggesting that clinical decisions regarding whether or not acupuncture should be recommended as a treatment option would be more accurately based on comparisons of acupuncture with usual care or acupuncture with other interventions (head-to-head studies and network meta-analyses) (25, 29). This trend of increased focus on pragmatic trials over explanatory trials is not limited to research into acupuncture, but has been identified as an issue in health research more generally (30). Further research on acupuncture mechanisms has also been recommended by both the Society for Acupuncture Research and the National Institutes for Health (NIH) in the USA which, at the Society for Acupuncture Research's biannual conference in Boston in October 2015, announced a new funding pool dedicated to acupuncture mechanism research (25). NIH have identified limited value in true versus sham acupuncture trials when investigating pain and have consigned such trials to a low programmatic priority (31). NIH has given high priority instead to pragmatic studies investigating acupuncture and pain management along with further research into the mechanisms of acupuncture (31).

The use of exit-debrief questionnaires in published sham/placebo-controlled acupuncture trials, which show no significant differences between the real and sham acupuncture groups in belief about which treatment they had received, suggests that in these trials placebo/nocebo effects are not an adequate explanation for any effects produced in the sham acupuncture group (32).

### **1.5 Acupuncture recommendations in clinical practice guidelines**

In Australia, acupuncture has been included in clinical practice guidelines for various types of acute pain including post-operative pain, and for rotator cuff syndrome (33, 34). In 'Acute Pain Management: Scientific Evidence' published by the Australian and New Zealand College of Anaesthetists and Faculty of Pain Medicine in 2015, NHMRC Level I evidence was identified from Cochrane reviews for acupuncture for labour pain, oocyte retrieval pain, primary dysmenorrhoea, tension-type headaches and migraine, and from PRISMA reviews for postoperative pain, back pain and acute burns pain (33).

*The development of guidelines requires quality levels of evidence adequate to support a given recommendation, even though they may fall below the highest level in some cases, and requires consideration of other factors including cost, clinical judgement, and patient preference.*

In 'Clinical Practice Guidelines for the Management of Rotator Cuff Syndrome in the Workplace', published by The University of New South Wales in 2013, Recommendation 23 states that 'Clinicians may consider acupuncture in conjunction with exercise; both modalities should be provided by suitably qualified health care providers' (Grade C: 'Body of evidence provides some support for recommendation but care should be taken in its application to individual clinical and organisational circumstances') (34).

In 'Consensus Guidelines for the Management of Postoperative Nausea and Vomiting' published in USA by the Society for Ambulatory Anesthesia in 2014, acupuncture and point stimulation of

PC6 were recommended as both prophylactic and treatment strategies [Category A recommendation: based on supportive literature which contains multiple randomised controlled trials which report statistically significant ( $P < 0.01$ ) differences between clinical interventions for a clinical outcome, and aggregated findings are supported by meta-analysis] (35).

Between 2012 and 2015 four International Symposia of Evidence-Based Clinical Practice Guideline in Traditional Medicine have been held in Daejeon, South Korea hosted by the Korean Institute of Oriental Medicine (KIOM), bringing together participants from Korea, China, Japan, UK, Norway, Holland and Australia (36). By November 2015, over 870 recommendations for acupuncture were identified for over 100 conditions from multiple international groups and over 30 countries (36). Examples include: the Otolaryngology Head Neck Surgery Foundation’s clinical practice guidelines for allergic rhinitis in 2015 (37); the National Institute for Health and Care Excellence (NICE), UK inclusion of migraine and tension type headache in 2012 (38); the Scottish Intercollegiate Guidelines Network guideline for the Management of Chronic Pain, inclusion of acupuncture (Grade A recommendation) for chronic low back pain and osteoarthritis (39); and the National German Gynaecologic Oncology Association’s (Arbeitsgemeinschaft Gynakologische Onkologie) inclusion of acupuncture for 12 symptoms associated with breast cancer treatment in 2015 (36, 37).

**Table 2. Statistical summary of findings of this review**

Evidence Level	Number of Conditions	Changes in Level of Evidence	Number of Conditions
Evidence of Positive effect	8	Increase to positive effect	5
Evidence of Potential positive effect	38	Increase to potential positive effect	18
Unclear/insufficient evidence	71	Increase to unclear/insufficient evidence	1
No evidence of effect	5	Decreased evidence level	2
Total conditions with some evidence of effect (any level)	117	_____	_____
Total conditions reviewed	122	Total increases in evidence level since prior reviews	24

### 1.6 Interpreting the findings

This review set out to identify the current state of evidence regarding acupuncture, and has done so principally by examining systematic reviews and meta-analyses. Most of these systematic reviews were restricted to only randomised controlled clinical trials which examine efficacy, not effectiveness (NHMRC level I evidence); however, some systematic reviews did include pragmatic trials and other uncontrolled studies. Consequently, studies which focused on effectiveness rather than efficacy such as cohort studies, case-control studies, or case series (NHMRC level III-IV evidence) were generally excluded from this review.

*Placebo controlled clinical trials consistently underestimate the true effect size of acupuncture, yet over 1000 trials have still demonstrated NHMRC Level I evidence for the effectiveness of acupuncture for 117 conditions.*

Earlier discussion has demonstrated the limitations of using placebo controlled clinical trials to assess acupuncture’s efficacy, with the NIH’s acknowledgement of this reflected in its current programmatic priorities which encourage

pragmatic trials at the expense of RCTs (31). Regardless of the limitations of acupuncture RCTs that under-report the true effect size of acupuncture (23, 28), for eight conditions across four clinical areas there is no longer any doubt of acupuncture's efficacy.

For the reasons outlined above, it is not correct to infer that acupuncture is ineffective for conditions which fall outside of the 'evidence of positive effect' category. For a further 109 conditions examined, positive results have been achieved from some trials in every case, with varying levels in the quality of the evidence.

Acupuncture has already been incorporated into clinical guidelines for over 100 conditions even though its current evidence level is rated below 'evidence of positive effect' in most of these cases. This reflects the weighting of other factors in guideline development, and recognition that the quality of levels of evidence are adequate to support a given recommendation, even though they may fall below the highest level in some cases (8). As recommended by both the Society for Acupuncture Research and the NIH, future acupuncture research to inform clinical practice guidelines should be focused on pragmatic trials, head-to-head comparison studies with other interventions (especially currently-recommended usual care interventions) to compare effectiveness, safety and cost-effectiveness, underpinned by further mechanism studies (25, 31). It is accurate to state from this report that there is NHMRC Level 1 evidence for acupuncture's effectiveness for 117 conditions.

The number of conditions included in this review is significantly larger than in previous reviews for two reasons. Firstly, acupuncture researchers have just begun to research the effectiveness of acupuncture for conditions not previously studied. Secondly research has become more nuanced and targeted. For example, what was previously reviewed as 'cancer adverse effects' is now separated into 12 different cancer-related conditions such as pain, fatigue, insomnia and xerostomia.

## **1.7 Limitations**

This review has several limitations. Literature before March 2013 was obtained by pooling results from the DVA 2010 and USVA 2013 reviews. Any MA or SR overlooked by those reviews were not identified in this review. MAs and SRs from March 2013 to September 2016 were identified via a PubMed search. MAs and SRs which were not indexed by PubMed were not identified in this review with one exception. Three additional SRs and MAs published between October 2016 and January 2017 were included in the final revision.

The findings of this review are limited by the mounting evidence that sham/placebo controls used in acupuncture trials are not inert, which is likely to lead to a consistent underestimation of the true effect size of acupuncture interventions (see sections 1.3 and 1.4 above).

By design this review aimed to update previous reviews, and focused on MAs and SRs. Studies which focus solely on effectiveness rather than efficacy such as cohort studies, case-control studies and case series were excluded by most of the SRs. This results in under-reporting studies on the effectiveness of acupuncture. For example, a PubMed search on 3 November 2016 for the following key words returned the following results:

- cohort AND acupuncture (208)
- case control AND acupuncture (377)
- case series AND acupuncture (150).

## **2 CONDITIONS NOW RATED AS ‘EVIDENCE OF POSITIVE EFFECT’ IN THIS REVIEW**

### **2.1 Migraine prophylaxis [Positive effect]**

For migraine prophylaxis, acupuncture was rated as ‘effective’ in the Australian DVA review (2010) and ‘evidence of positive effect’ in the USVA Evidence map of acupuncture (2014) (5, 6). Since March 2013 a narrative review of high quality randomised controlled trials and two systematic reviews including a Cochrane systematic review update, have confirmed that acupuncture is superior to sham acupuncture and seems to be at least as effective as conventional preventative medication in reducing migraine frequency (40-42). Moreover, acupuncture is described as ‘safe, long-lasting and cost effective’ (40). Subgroup analysis in the Cochrane systematic review found that 16 or more treatment sessions showed a larger effect size ( $Z=4.06$ ) than 12 treatments or fewer ( $Z=2.32$ ). Evidence levels in these three reviews was moderate to high quality.

### **2.2 Headache (chronic tension-type and chronic episodic) [Positive effect]**

Chronic tension-type headaches and chronic episodic headaches were not reviewed in the Australian DVA review (2010) and rated as ‘evidence of positive effect’ in the USVA Evidence map of acupuncture (2014) (5, 6). The most recent Cochrane systematic review update confirmed that acupuncture is effective for frequent episodic and chronic tension-type headaches with moderate to low quality evidence (43). A brief review of systematic reviews and meta-analyses described acupuncture as having a ‘potentially important role as part of a treatment plan for migraine, tension-type headache, and several different types of chronic headache disorders’ (44). Studies in Germany and the UK found acupuncture for chronic headaches to be cost-effective (44).

### **2.3 Low back pain [Chronic – positive effect; acute – potential positive effect]**

For low back pain, acupuncture was rated as ‘effective (possibly)’ in the Australian DVA review (2010) and ‘unclear’ in the USVA Evidence map of acupuncture (2014) (5, 6). The main reason given for the ‘unclear’ rating in the USVA Evidence map was that ‘sham acupuncture controlled trials tended towards statistically nonsignificant results’ and a Cochrane systematic review in 2005 which drew no firm conclusions on the effectiveness of acupuncture for acute low back pain (6). The problems associated with the interpretation of the effects of sham acupuncture have already been discussed. When reviews separate the evidence on chronic low back pain from acute low back pain, as was done in the recent review by the US Agency for Healthcare Research and Quality, it was found that there is moderate quality evidence for the effectiveness of acupuncture in chronic low back pain for both pain intensity and function, but only low quality evidence for pain intensity and function in acute low back pain (45).

In a systematic review of 32 randomised controlled trials for acupuncture treatment of chronic non-specific low back pain, acupuncture was superior (both statistically and clinically significant) to sham acupuncture in pain reduction and improved function immediately after treatment (46). Acupuncture was equivalent but not superior to usual care in pain and function, but acupuncture plus usual care was superior to usual care alone (46). The reviewers rated the evidence level of their review as Level of Evidence I (46). A systematic review of 11 randomised controlled trials on acute low back pain, acupuncture was superior to NSAIDs for improving symptoms (small effect), and superior to sham for pain but not function (47). A narrative review of non-invasive and alternative treatments for chronic low back pain rated the evidence for the effectiveness of acupuncture as high and for acupressure as moderate (48).

A review of 16 systematic reviews found that acupuncture alone, or when added to usual care, provided short-term improvement in pain and function for chronic low back pain (medium to large clinical effects) and hence 'should be advocated in routine clinical practice' (49). For acute low back pain, the reviewers could not make firm conclusions about the effectiveness of acupuncture due to the inclusion of only two systematic reviews (49). Two studies found that acupuncture is likely to be cost-effective for low back pain or chronic non-specific low back pain, respectively (50, 51).

#### **2.4 Knee osteoarthritis pain [Positive effect]**

Knee osteoarthritis pain was not reviewed in the Australian DVA review (2010) and rated as 'evidence of potential positive effect' in the USVA Evidence map of acupuncture (2014) (5, 6). In a network meta-analysis comparing 22 interventions in 152 studies, acupuncture was found to be equal to balneotherapy and superior to sham acupuncture, muscle-strengthening exercise, Tai Chi, weight loss, standard care and aerobic exercise (in ranked order) (52). Acupuncture was also superior to standard care and muscle-strengthening exercises in a sub-analysis of moderate to high quality studies (52). In a systematic review of 12 randomised controlled trials, acupuncture was found to significantly reduce pain intensity, to improve functional mobility and quality of life (53). Subgroup analysis showed greater reduction in pain intensity when treatment lasted for more than four weeks (53). The reviewers concluded that 'current evidence supports the use of acupuncture as an alternative for traditional analgesics in patients with osteoarthritis' (53).

#### **2.5 Allergic rhinitis (seasonal and perennial/persistent) [Positive effect]**

For allergic rhinitis, acupuncture was rated as 'effective' in the Australian DVA review (2010) and 'unclear' in the USVA Evidence map of acupuncture (2014) (5, 6). A systematic review of 13 randomised controlled trials concluded that 'acupuncture could be a safe and valid treatment option for allergic rhinitis' (moderate quality evidence) (54). Another systematic review (which included two large multi-centre randomised controlled trials, three comparisons of acupuncture versus medication and one cost-effectiveness study) concluded that there is high quality evidence of the efficacy and effectiveness of acupuncture and that it appears to be safe and cost-effective (15). Clinical practice guidelines for allergic rhinitis published by the Otolaryngology Head Neck Surgery Foundation in 2015 included acupuncture as Option five: 'Clinicians may offer acupuncture, or refer to a clinician who can offer acupuncture, for patients with AR who are interested in nonpharmacological therapy' (Aggregate evidence quality - Grade B) (37).

#### **2.6 Chemotherapy-induced nausea and vomiting (CINV) [Positive effect]**

For chemotherapy-induced nausea and vomiting, acupuncture was rated as 'effective' in the Australian DVA review (2010) and was not reviewed separately to 'cancer adverse effects' in the USVA Evidence map of acupuncture (2014) (5, 6). In 2013, a systematic review of seven acupuncture and six acupressure RCTs found that acupuncture reduced the frequency of acute vomiting and the dose of rescue medication but did not reduce acute nausea severity or frequency compared to control. Acupressure showed a decrease in frequency of nausea but not acute vomiting or delayed symptoms. All studies used state-of-the-art combination anti-emetics in addition to acupuncture/acupressure. The reviewers rated the quality of evidence in the acupuncture studies as low to moderate while the acupressure studies were moderate to high quality. There was insufficient evidence to draw firm conclusions due to underpowered studies

(21). However, an updated systematic review by Garcia et al in 2014 (using 18 new RCTs) found that acupuncture is an appropriate referral option for chemotherapy-induced nausea and vomiting (55). On this basis, CINV has been rated as positive rather than potential positive in this review.

## **2.7 Post-operative nausea and vomiting (PONV) [Positive effect]**

For post-operative nausea and vomiting (PONV), acupuncture was rated as ‘insufficient evidence’ in the Australian DVA review (2010) and ‘potential positive effect’ in the USVA Evidence map of acupuncture (2014) (5, 6). In a systematic review and meta-analysis of 30 RCTs on acupuncture and acupressure in 2013, both acupuncture and acupressure reduced the number of cases of early nausea and vomiting (up to 24 hours post-surgery) (56). In 2015, a Cochrane update of 59 RCTs on PC6 stimulation found that PC6 stimulation was superior to sham, and equivalent to modern anti-emetics (16).

## **2.8 Post-operative pain [Positive effect]**

For post-operative pain, acupuncture was not reviewed in the Australian DVA review (2010) and rated as ‘unclear’ in the USVA Evidence map of acupuncture (2014) (5, 6). A systematic review and meta-analysis of 13 RCTs in 2016, found that acupuncture, electroacupuncture (EA) and transcutaneous electrical acupoint stimulation (TEAS) improved pain on day one after surgery and reduced opioid use (17). Subgroup analysis showed that acupuncture and TEAS were superior to EA (17). A systematic review specifically on acute pain after back surgery reviewed five RCTs (three of which were high quality) and found encouraging but limited evidence for the efficacy of acupuncture (57). A systematic review on complementary therapies for pain after knee surgery included three RCTs on acupuncture and one on acupressure (58). The review found that acupressure reduced pain, and while acupuncture did not reduce pain it did reduce ibuprofen use (58). An RCT on acupuncture for pain after total knee arthroplasty found that acupuncture was superior to sham in post-operative fentanyl use, time to first request for fentanyl and pain intensity (59).

### 3 RESEARCH INTO THE MECHANISMS OF ACUPUNCTURE

Mechanisms underlying acupuncture analgesia have been extensively researched for over 60 years. In animal models, acupuncture and/or electroacupuncture have been shown to be effective for the alleviation of inflammatory, neuropathic, cancer, and visceral pain (60). Ascending neural pathways involving A $\delta$ , A $\beta$  and C sensory fibres have been mapped, the mesolimbic loop of analgesia in the brain and brain stem has been identified and descending pathways have also been mapped (61). Numerous mediators have been identified including opioid and non-opioid neuropeptides, serotonin, norepinephrine, dopamine, cytokines, glutamate, nitric oxide and gamma-amino-butyric-acid (GABA) (60, 61). Acupuncture analgesia has been shown to involve several classes of opioid neuropeptides including enkephalins, endorphins, dynorphins, endomorphins and nociceptin (also known as Orphanin FQ) (61-63). Among the non-opioid neuropeptides, substance P (SP), vasoactive intestinal peptide (VIP) and calcitonin gene-related peptide (CGRP) have been investigated for their roles in both the analgesic and anti-inflammatory effects of acupuncture (60, 64). Two recent reviews of acupuncture analgesia research further demonstrate the complexity of this area of study (61, 62).

The anti-inflammatory effects of acupuncture involve numerous mediators, receptors and signalling pathways, as outlined in two recent reviews (64, 65). The anti-inflammatory effects of acupuncture have particular relevance to allergic rhinitis, irritable bowel syndrome, post-surgical recovery, migraine, osteoarthritis and inflammatory aspects of a range of musculoskeletal conditions. In allergic rhinitis, acupuncture has been shown to down-regulate total and specific IgE, as well as SP and VIP (32, 66). Acupuncture has been shown to down-regulate transient receptor potential vanilloid 1 (TRPV1) in inflammatory pain and there is indirect evidence to suggest that acupuncture may down-regulate TRPV1 expression and sensitivity in allergic rhinitis (32, 64, 67). In irritable bowel syndrome, acupuncture has been shown to down-regulate SP, VIP and CGRP (68, 69). In migraine, acupuncture has been reported to down-regulate CGRP and SP which are also powerful vasodilators (70, 71).

In addition to the extensive research literature on acupuncture's efficacy, effectiveness and safety there is a body of research (largely using animal models) which has investigated physiological changes underpinning the effects of acupuncture in a broad range of clinical areas apart from pain and inflammation. To canvas this research in detail is beyond the scope of this review, however numerous reviews of this mechanism research have been published. A PubMed search on 18 September 2016, using the search terms 'acupuncture AND mechanism', yielded 1,943 hits.

For example:

- Acupuncture regulation of female reproductive function (72)
- Acupuncture regulation of gastrointestinal function (73)
- Acupuncture regulation of bladder function (73)
- Acupuncture regulation of circulation (74).

**Table 3. Summary of effectiveness/efficacy in acupuncture research literature sorted by evidence levels**

Australian DVA (Sept 2005 - Sept 2010)	USVA Evidence map (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)	
Effective	Evidence of positive effect	Evidence of positive effect	
<ul style="list-style-type: none"> <li>- Allergic rhinitis (for up to 3 months)</li> <li>- Back or pelvic pain during pregnancy</li> <li>- Migraine prophylaxis (when used with routine care)</li> <li>- Vomiting after chemotherapy (with superseded anti-emetic drugs)</li> </ul>	<ul style="list-style-type: none"> <li>- Chronic pain*</li> <li>- Headache</li> <li>- Migraine</li> </ul> <p><i>* Chronic pain has been separated into different chronic pain conditions since USVA</i></p>	<ul style="list-style-type: none"> <li>- Allergic rhinitis (perennial &amp; seasonal)</li> <li>- Chemotherapy-induced nausea and vomiting (CINV) (with anti-emetics)</li> <li>- Chronic low back pain</li> <li>- Headache (tension-type and chronic)</li> <li>- Knee osteoarthritis</li> <li>- Migraine prophylaxis</li> <li>- Postoperative nausea &amp; vomiting</li> <li>- Postoperative pain</li> </ul>	
Effective (possibly)	Evidence of potential positive effect		
<ul style="list-style-type: none"> <li>- Lower back pain</li> </ul>	<ul style="list-style-type: none"> <li>- Ankle sprain</li> <li>- Anxiety</li> <li>- Cancer pain</li> <li>- Constipation</li> <li>- Depression</li> <li>- Dysmenorrhoea</li> <li>- General pain</li> <li>- Insomnia</li> <li>- Labour pain</li> <li>- Obesity</li> <li>- Osteoarthritic pain</li> <li>- Plantar heel pain</li> <li>- Postoperative nausea and vomiting</li> <li>- Post-traumatic Stress Disorder</li> <li>- Pregnancy pain</li> <li>- Prostatitis pain</li> <li>- Restless leg syndrome</li> <li>- Schizophrenia</li> <li>- Smoking cessation</li> <li>- Temporomandibular pain</li> </ul>	<ul style="list-style-type: none"> <li>- Acute low back pain</li> <li>- Acute stroke</li> <li>- Ambulatory anaesthesia</li> <li>- Anxiety</li> <li>- Aromatase-inhibitor-induced arthralgia</li> <li>- Asthma in adults</li> <li>- Back or pelvic pain during pregnancy</li> <li>- Cancer pain</li> <li>- Cancer-related fatigue</li> <li>- Constipation</li> <li>- Craniotomy anaesthesia</li> <li>- Depression (with antidepressants)</li> <li>- Dry eye</li> <li>- Hypertension (with medication)</li> <li>- Insomnia</li> <li>- Irritable bowel syndrome</li> <li>- Labour pain</li> <li>- Lateral elbow pain</li> <li>- Menopausal hot flushes</li> <li>- Modulating sensory perception thresholds</li> <li>- Neck pain (NAD, not WAD)</li> </ul>	<ul style="list-style-type: none"> <li>- Obesity</li> <li>- Perimenopausal &amp; postmenopausal insomnia</li> <li>- Plantar heel pain</li> <li>- Post-stroke insomnia</li> <li>- Post-stroke shoulder pain</li> <li>- Post-stroke spasticity</li> <li>- Post-traumatic stress disorder</li> <li>- Prostatitis pain/chronic pelvic pain syndrome</li> <li>- Recovery after colorectal cancer resection</li> <li>- Restless leg syndrome</li> <li>- Schizophrenia (with antipsychotics)</li> <li>- Sciatica</li> <li>- Shoulder impingement syndrome (early stage) (with exercise)</li> <li>- Shoulder pain</li> <li>- Smoking cessation (up to 3 months)</li> <li>- Stroke rehabilitation</li> <li>- Temporomandibular pain</li> </ul>

**Table 3. Summary of effectiveness/efficacy in acupuncture research literature sorted by evidence levels (contd)**

Australian DVA (Sept 2005 - Sept 2010)	USVA Evidence map (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)	
Insufficient evidence	Unclear evidence	Unclear/insufficient evidence	
<ul style="list-style-type: none"> <li>- Assisted conception (when used on the day of embryo transfer)</li> <li>- Asthma</li> <li>- Bell's palsy</li> <li>- Breathlessness</li> <li>- Chronic neck pain</li> <li>- Cocaine addiction</li> <li>- Depression</li> <li>- Glaucoma</li> <li>- Induction of labour</li> <li>- Insomnia</li> <li>- Irritable bowel syndrome</li> <li>- Labour pain</li> <li>- Lateral elbow pain</li> <li>- Postnatal depression</li> <li>- Postoperative nausea and vomiting</li> <li>- Restless leg syndrome</li> <li>- Rheumatoid arthritis</li> <li>- Schizophrenia</li> <li>- Shoulder pain</li> <li>- Smoking cessation (up to 3 months)</li> <li>- Stroke</li> <li>- Uterine fibroids</li> <li>- Vascular dementia</li> </ul>	<ul style="list-style-type: none"> <li>- Back pain</li> <li>- Cancer adverse effects</li> <li>- Chronic fatigue syndrome</li> <li>- Drug addiction</li> <li>- Dry eye</li> <li>- Erectile dysfunction</li> <li>- Exercise</li> <li>- Fibromyalgia pain</li> <li>- Gastrointestinal disease</li> <li>- High blood pressure</li> <li>- Irritable bowel syndrome</li> <li>- Menopausal symptoms</li> <li>- Neck pain</li> <li>- Opiate addiction</li> <li>- Postoperative pain</li> <li>- Premenstrual syndrome</li> <li>- Quality of life</li> <li>- Rheumatoid arthritis pain</li> <li>- Rhinitis</li> <li>- Shoulder pain</li> <li>- Surgical analgesia</li> <li>- Tinnitus</li> <li>- Xerostomia</li> </ul>	<ul style="list-style-type: none"> <li>- Acupuncture in Emergency Department</li> <li>- Acute ankle sprain in adults</li> <li>- Alzheimer's disease</li> <li>- Angina pectoris</li> <li>- Assisted conception in ART (includes SR and MA from Dec 2016 and Jan 2017)</li> <li>- Asthma in children</li> <li>- Atopic dermatitis</li> <li>- Attention Deficit Hyperactivity Disorder (ADHD)</li> <li>- Autism spectrum disorder (ASD)</li> <li>- Bell's palsy</li> <li>- Bladder pain syndrome</li> <li>- Cancer-related insomnia</li> <li>- Cancer-related psychological symptoms</li> <li>- Carpal tunnel syndrome</li> <li>- Chemotherapy-induced peripheral neuropathy</li> <li>- Chronic fatigue syndrome</li> <li>- Chronic kidney disease</li> <li>- Chronic obstructive pulmonary disease (COPD)</li> <li>- Chronic urinary retention due to spinal cord injury</li> <li>- Chronic urticaria</li> </ul>	<ul style="list-style-type: none"> <li>- Dysmenorrhoea</li> <li>- Dyspepsia in diabetic gastroparesis (DGP)</li> <li>- Erectile dysfunction</li> <li>- Exercise performance &amp; post-exercise recovery</li> <li>- Fatigue in systemic lupus erythematosus</li> <li>- Fibromyalgia</li> <li>- Functional dyspepsia</li> <li>- Gag reflex in dentistry</li> <li>- Glaucoma</li> <li>- Heart failure</li> <li>- Hot flushes in breast cancer</li> <li>- Hyperemesis gravidarum</li> <li>- Hypoxic ischemic encephalopathy in neonates</li> <li>- Induction of labour</li> <li>- Inflammatory bowel disease</li> <li>- Itch</li> <li>- Lumbar spinal stenosis</li> <li>- Melasma</li> <li>- Meniere's disease/syndrome</li> <li>- Menopausal syndrome</li> <li>- Multiple sclerosis</li> <li>- Mumps in children</li> <li>- Myelosuppression after chemotherapy</li> <li>- Oocyte retrieval pain relief</li> </ul>

**Table 3. Summary of effectiveness/efficacy in acupuncture research literature sorted by evidence levels (contd)**

Australian DVA (Sept 2005 - Sept 2010)	USVA Evidence map (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
<b>Insufficient evidence</b>	<b>Unclear evidence</b>	<b>Unclear/insufficient evidence</b>
		<ul style="list-style-type: none"> <li>- Opiate addiction</li> <li>- Opioid detoxification</li> <li>- Parkinson’s disease</li> <li>- Polycystic ovarian syndrome</li> <li>- Poor sperm quality</li> <li>- Postnatal depression</li> <li>- Postoperative gastroparesis syndrome (PGS)</li> <li>- Postoperative ileus</li> <li>- Post-stroke hiccoughs</li> <li>- Premenstrual syndrome</li> <li>- Primary ovarian insufficiency</li> <li>- Primary Sjogren’s syndrome</li> <li>- Psoriasis vulgaris</li> <li>- Rheumatoid arthritis</li> <li>- Slowing progression of myopia</li> <li>- Spinal cord injury</li> <li>- Stress urinary incontinence in adults</li> <li>- Sudden sensorineural hearing loss</li> <li>- Surgery analgesia</li> <li>- Tinnitus</li> <li>- Traumatic brain injury</li> <li>- Urinary incontinence</li> <li>- Uterine fibroids</li> <li>- Vascular cognitive impairment without dementia</li> <li>- Vascular dementia</li> <li>- Whiplash associated disorder (WAD)</li> <li>- Xerostomia in cancer</li> </ul>
<b>Not effective</b>	<b>Evidence of no effect</b>	<b>No evidence of effect</b>
<ul style="list-style-type: none"> <li>- Assisted conception (if used around time of oocyte retrieval)</li> <li>- Epilepsy</li> <li>- Smoking cessation (more than 3 months)</li> </ul>	<ul style="list-style-type: none"> <li>- Alcohol dependence</li> <li>- Carpal tunnel syndrome pain</li> <li>- Cocaine addiction</li> <li>- Nausea in pregnancy</li> </ul>	<ul style="list-style-type: none"> <li>- Alcohol dependence</li> <li>- Cocaine addiction</li> <li>- Epilepsy</li> <li>- Nausea in pregnancy</li> <li>- Smoking cessation (more than 6 months)</li> </ul>

**Table 4. Summary of changes in evidence levels from 2005 to 2016**

Conditions which have not changed in evidence level	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
Migraine prophylaxis	Effective	<b>Positive effect</b>	<b>Positive effect</b>
Headache (tension-type and chronic)	Not reviewed	<b>Positive effect</b>	<b>Positive effect</b>
Chemotherapy-induced nausea and vomiting	<b>Effective</b>	Not reviewed	<b>Positive effect</b>
Allergic rhinitis	<b>Effective</b>	Unclear	<b>Positive effect</b>
Anxiety	Not reviewed	<b>Potential positive effect</b>	<b>Potential positive effect</b>
Post-traumatic Stress Disorder	Not reviewed	<b>Potential positive effect</b>	<b>Potential positive effect</b>
Schizophrenia	Insufficient evidence	<b>Potential positive effect</b>	<b>Potential positive effect</b>
Smoking cessation		<b>Potential positive effect</b>	
Up to 3 months post-treatment	Insufficient evidence		<b>Potential positive effect</b>
More than 3 months post-treatment	Not effective		<b>No evidence of effect (more than 6 months)</b>
Temporomandibular pain		<b>Potential positive effect</b>	<b>Potential positive effect</b>
Plantar heel pain		<b>Potential positive effect</b>	<b>Potential positive effect</b>
Cancer pain	Not reviewed	<b>Potential positive effect</b>	<b>Potential positive effect</b>
Depression	Insufficient evidence	<b>Potential positive effect</b>	<b>Potential positive effect</b>
Insomnia	Insufficient evidence	<b>Potential positive effect</b>	<b>Potential positive effect</b>
Labour pain	Insufficient evidence	<b>Potential positive effect</b>	<b>Potential positive effect</b>
Back or pelvic pain during pregnancy	Effective	<b>Potential positive effect</b>	<b>Potential positive effect</b>
Prostatitis pain/chronic pelvic pain syndrome	Not reviewed	<b>Potential positive effect</b>	<b>Potential positive effect</b>
Constipation	Not reviewed	<b>Potential positive effect</b>	<b>Potential positive effect</b>
Obesity	Not reviewed	<b>Potential positive effect</b>	<b>Potential positive effect</b>
Restless leg syndrome	Insufficient evidence	<b>Potential positive effect</b>	<b>Potential positive effect</b>
Induction of labour	<b>Insufficient evidence</b>	Not reviewed	Unclear
Bell's palsy	<b>Insufficient evidence</b>	Not reviewed	Unclear
Glaucoma	<b>Insufficient evidence</b>	Not reviewed	Unclear
Uterine fibroids	<b>Insufficient evidence</b>	Not reviewed	Unclear
Vascular dementia	<b>Insufficient evidence</b>	Not reviewed	Unclear
Erectile dysfunction	Not reviewed	Unclear	Unclear
Tinnitus	Not reviewed	Unclear	Unclear
Improving exercise performance/recovery	Not reviewed	Unclear	Unclear
Opiate addiction	Not reviewed	Unclear	Unclear
Chronic fatigue syndrome	Not reviewed	Unclear	Unclear

**Table 4. Summary of changes in evidence levels from 2005 to 2016 (contd)**

Conditions which have not changed in evidence level	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
<b>Fibromyalgia</b>	Not reviewed	<b>Unclear</b>	<b>Unclear</b>
<b>Premenstrual syndrome</b>	Not reviewed	<b>Unclear</b>	<b>Unclear</b>
<b>Rheumatoid arthritis</b>	Insufficient evidence	<b>Unclear</b>	<b>Unclear</b>
<b>Assisted conception in ART</b>	<b>Insufficient evidence</b> at embryo transfer; <b>Ineffective</b> at oocyte retrieval	Not reviewed	<b>Unclear</b>
<b>Nausea in pregnancy</b>	Not reviewed	<b>Evidence of no effect</b>	<b>No evidence of effect</b>
<b>Alcohol dependence</b>	Not reviewed	<b>Evidence of no effect</b>	<b>No evidence of effect</b>
<b>Cocaine addiction</b>	Insufficient evidence	<b>Evidence of no effect</b>	<b>No evidence of effect</b>
<b>Epilepsy</b>	<b>Not effective</b>	Not reviewed	<b>No evidence of effect</b>
Conditions which have increased in evidence level	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
<b>Low back pain</b>	Possibly effective	Unclear	<i>Research has shown better outcomes for chronic low back pain than for acute low back pain</i>
Chronic low back pain			<b>Positive effect</b>
Acute low back pain			<b>Potential positive effect</b>
<b>Knee osteoarthritis</b>	Not reviewed	<b>Potential positive effect</b>	<b>Positive effect</b>
<b>Postoperative nausea and vomiting</b>	Insufficient evidence	<b>Potential positive effect</b>	<b>Positive effect</b>
<b>Postoperative pain</b>	Not reviewed	<b>Unclear</b>	<b>Positive effect</b>
Postoperative nausea and vomiting and pain after tonsillectomy			<b>Positive effect</b>
Postoperative pain – back surgery			<b>Positive effect</b>
Postoperative pain – knee surgery			<b>Positive effect</b>
<b>Stroke</b>	<b>Insufficient evidence</b>	Not reviewed	<i>Stroke research has now separated into several new topics</i>
Acute stroke			<b>Potential positive effect</b>
Stroke rehabilitation			<b>Potential positive effect</b>
Post-stroke spasticity			<b>Potential positive effect</b>
Post-stroke insomnia			<b>Potential positive effect</b>
Post-stroke shoulder pain			<b>Potential positive effect</b>
Post-stroke hiccoughs			<b>Unclear</b>

**Table 4. Summary of changes in evidence levels from 2005 to 2016 (contd)**

Conditions which have increased in evidence level	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
<b>Menopausal symptoms</b>	Not reviewed	<b>Unclear</b>	<i>Menopausal symptom research has now separated into two new topics</i>
Menopausal hot flushes			<b>Potential positive effect</b>
Perimenopausal and postmenopausal sleep disturbance			<b>Potential positive effect</b>
<b>Cancer adverse effects</b> (not including cancer pain and chemotherapy-induced nausea and vomiting)	Not reviewed	<b>Unclear</b>	<i>Cancer adverse effects research has now separated into several new topics</i>
Cancer-related fatigue			<b>Potential positive effect</b>
Cancer-related insomnia			<b>Unclear</b>
Cancer-related peripheral neuropathy			<b>Unclear</b>
Hot flushes/flushes in breast cancer			<b>Unclear</b>
Xerostomia in cancer	Not reviewed	<b>Unclear</b>	<b>Unclear</b>
Recovery after colorectal cancer resection			<b>Potential positive effect</b>
Aromatase-inhibitor-induced arthralgia			<b>Potential positive effect</b>
Chemotherapy-induced peripheral neuropathy			<b>Unclear</b>
Myelosuppression after chemotherapy			<b>Unclear</b>
Cancer-related psychological symptoms			<b>Unclear</b>
<b>Surgical analgesia</b>	Not reviewed	<b>Unclear</b>	<i>Surgical analgesia research has now separated into several new topics</i>
Craniotomy anaesthesia			<b>Potential positive effect</b>
Ambulatory anaesthesia			<b>Potential positive effect</b>
<b>Dry eye</b>	Not reviewed	<b>Unclear</b>	<b>Potential positive effect</b>
<b>Irritable bowel syndrome</b>	Insufficient evidence	<b>Unclear</b>	<b>Potential positive effect</b>
<b>Hypertension</b>	Not reviewed	<b>Unclear</b>	<b>Potential positive effect</b>
<b>Lateral elbow pain</b>	<b>Insufficient evidence</b>	Not reviewed	<b>Potential positive effect</b>

<b>Table 4. Summary of changes in evidence levels from 2005 to 2016 (contd)</b>			
<b>Conditions which have increased in evidence level</b>	<b>Australian DVA (Sept 2005 - Sept 2010)</b>	<b>USVA (Jan 2005 - Mar 2013)</b>	<b>The Acupuncture Evidence Project (Mar 2013 - Sept 2016)</b>
	<i>Baseline evidence in bold</i>		
<b>Neck pain</b>			
Neck pain and associated disorders (NAD)	Insufficient evidence	<b>Unclear</b>	<b>Potential positive effect</b>
Whiplash associated disorders (WAD)			<b>Unclear</b>
<b>Shoulder pain</b>	Insufficient evidence	<b>Unclear</b>	<b>Potential positive effect</b>
<b>Asthma</b>	<b>Insufficient evidence</b>	Not reviewed	<b>Potential positive effect (in adults)</b>
<b>Carpal tunnel syndrome</b>	Not reviewed	<b>Evidence of no effect</b>	<b>Unclear</b>
<b>Conditions which have decreased in evidence level</b>	<b>Australian DVA (Sept 2005 - Sept 2010)</b>	<b>USVA (Jan 2005 - Mar 2013) (to Mar 2013)</b>	<b>The Acupuncture Evidence Project (to September 2016)</b>
	<i>Baseline evidence in bold</i>		
<b>Dysmenorrhoea</b>	Not reviewed	<b>Potential positive effect</b>	<b>Unclear</b>
<b>Ankle sprain</b>	Not reviewed	<b>Potential positive effect</b>	<b>Unclear</b>

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references**

Conditions which remain unchanged	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
<b>Migraine prophylaxis</b>	Effective	<b>Positive effect</b>	<p>Acupuncture seems to be at least as effective as conventional preventative medication for migraine and is safe, long lasting, and cost-effective (citing Witt et al 2008) (Da Silva 2015 - Narrative review of large high quality RCTs)(40)</p> <p>Acupuncture reduces migraine frequency. Acupuncture superior to sham and may be as effective as prophylactic drugs; subgroup analysis showed larger effect size when 16 or more treatments given (<math>Z = 4.06</math>) compared to 12 treatments or less (<math>Z = 2.32</math>); Moderate quality evidence (Linde 2016 - Cochrane SR of 22 RCTs of at least 8 weeks duration)(41)</p> <p>Acupuncture superior to sham in effectiveness and reduced risk of recurrence; Moderate to high quality evidence (Yang 2016 - SR of 10 RCTs)(42)</p>
<b>Chemotherapy-induced nausea and vomiting</b>	<b>Effective</b>	Not reviewed	<p>Acupuncture reduced the frequency of acute vomiting and the dose of rescue medication but did not reduce acute nausea severity or frequency compared to control. Acupressure showed a decrease in frequency of nausea but not acute vomiting or delayed symptoms. All studies used state-of-the-art combination anti-emetics. Insufficient evidence due to underpowered studies; acupuncture low to moderate quality evidence; acupressure moderate to high quality evidence (McKeon 2013 - SR of 7 acupuncture and 6 acupressure RCTs)(21)</p> <p>Acupuncture is an appropriate referral option for chemotherapy-induced nausea and vomiting (Garcia 2014 - SR update: 18 new RCTs)(55)</p>
<b>Allergic rhinitis</b>	<b>Effective</b>	Unclear	<p>Acupuncture could be a safe and valid treatment for allergic rhinitis; Moderate quality evidence (Feng 2015 - SR of 13 RCTs)(54)</p> <p>High quality evidence of efficacy and effectiveness (Taw 2015 - SR of 2 large multi-centre RCTs, 3 acupuncture vs medication RCTs and 1 cost-effectiveness study)(15)</p> <p>OHNSF clinical practice guideline: Option 5: Clinicians may offer acupuncture, or refer to a clinician who can offer acupuncture, for patients with AR who are interested in nonpharmacologic therapy; Aggregate evidence quality - Grade B (Seidman 2015)(37)</p> <p>Acupuncture is cost-effective for allergic rhinitis (Kim 2012, Witt 2010)(75, 76)</p> <p>SAR - Acupuncture significantly superior to rescue medication in QALY gained, but may cost more short term (Reinhold 2013)(77, 78)</p>

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which remain unchanged	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
<b>Anxiety</b>	Not reviewed	<b>Potential positive effect</b>	Acupuncture has ‘potential use’ (Bazzan 2014 - Narrative review)(79) Positive and statistically significant effects; Moderate to high quality evidence (Goyata 2016 - Integrative review of 19 studies including 6 RCTs; 11 high quality studies; 5 moderate)(80)
<b>Post-traumatic Stress Disorder</b>	Not reviewed	<b>Potential positive effect</b>	One new RCT - Acupuncture plus usual care superior to usual care in PTSD severity, depression, pain and physical and mental functioning (Engel 2015 - RCT acupuncture plus usual care vs usual care, n = 55)(81)
<b>Schizophrenia</b>	Insufficient evidence	<b>Potential positive effect</b>	Acupuncture effective for schizophrenia, especially in improving sleep, mood and QoL by modulating and normalizing the limbic–paralimbic–neocortical network (LPNN), including the default mode network (DMN); limited evidence (Bosch 2015 - Review of SRs & MAs)(82) Acupuncture plus antipsychotic medication superior to antipsychotic medication alone, in terms of mental state and length of hospitalisation (moderate quality evidence) with fewer adverse effects (low quality evidence) (Shen 2014 – Cochrane update of 30 RCTs)(83)
<b>Smoking cessation</b>	Insufficient evidence (up to 3 months); Not effective (more than 3 months)	<b>Potential positive effect (short term)</b>	Potential short-term effects (low quality evidence), Insufficient evidence (more than 6 months) (White 2014 – Cochrane update)(84)
<b>Temperomandibular pain</b>		<b>Potential positive effect</b>	One new RCT: Acupuncture was equivalent to occlusal splint in pain intensity and range of mouth opening (Grillo 2015 - RCT acupuncture vs occlusal splint, n = 40)(85)
<b>Plantar heel pain</b>		<b>Potential positive effect</b>	No updates
<b>Cancer pain</b>	Not reviewed	<b>Potential positive effect</b>	Conflicting SRs Insufficient evidence; low to very low quality evidence (Paley 2015 - Cochrane SR of 5 RCTs) (86) Acupuncture relieved malignancy-related and surgery-induced pain but not pain induced by chemotherapy, radiotherapy or hormone therapy; Reviewers recommend acupuncture be included in multimodal treatment regimens (Chiu 2016 - SR of 29 RCTs)(87)

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which remain unchanged	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
<b>Depression</b>	Insufficient evidence	<b>Potential positive effect</b>	Acupuncture plus SSRIs superior to SSRIs alone, with an early onset of action and was safe and well-tolerated; EA had greater effect than manual acupuncture (Chan 2015 – SR of 13 RCTs: 1 high quality trial, 5 moderate, 7 low) (88) Effective and safe for major depressive disorder, especially in improving sleep, mood and QoL by modulating and normalizing the limbic–paralimbic–neocortical network (LPNN), including the default mode network (DMN); ‘promising evidence’ (Bosch 2015 - Review of SRs & MAs)(82) Acupuncture is cost-effective compared with counselling or usual care alone, although the ranking of counselling and acupuncture depends on the relative cost of delivering these interventions (Spackman 2014 –cost-effectiveness in one RCT)(89)
<b>Insomnia</b>	Insufficient evidence	<b>Potential positive effect</b>	Acupuncture may be superior to medication. Acupuncture for insomnia is potentially mediated by norepinephrine, melatonin, gamma-aminobutyric acid, and beta-endorphin; insufficient evidence (Zhao 2013 - book chapter)(90) Acupuncture statistically superior to sham (3 studies) and medication (27 studies); low quality evidence (Shergis 2016 - SR)(91)
<b>Labour pain</b>	Insufficient evidence	<b>Potential positive effect</b>	Acupuncture & acupressure ‘promising’ – Conflicting results due to heterogeneity in study designs, research questions, treatment protocols and outcomes measures (Levett 2014 - Review of SRs)(92)
<b>Back or pelvic pain during pregnancy</b>	Effective	<b>Potential positive effect</b>	Promising results; low quality evidence (Selva Olid 2013)(93) Clinically important and statistically significant changes (Close 2014 - SR of CAM; 2 high quality studies)(94) Moderate quality evidence that acupuncture or exercise, tailored to the stage of pregnancy, significantly reduced evening pelvic or lumbo-pelvic pain. Acupuncture superior to exercise for reducing evening pelvic pain; Both acupuncture and exercise superior to usual care; Insufficient evidence (Liddle 2015 - Cochrane SR: Comparison of interventions)(95)

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which remain unchanged	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
<b>Prostatitis pain/chronic pelvic pain syndrome</b>	Not reviewed	<b>Potential positive effect</b>	Acupuncture superior to both sham and to usual care and safe, thus it should be offered when available (Chang 2016 - SR of 7 RCTs: 3 high quality studies, 1 moderate and 3 low)(96) Acupuncture superior to sham in pain, voiding and QoL; acupuncture superior to medication in pain relief but no different in voiding and QoL; acupuncture as an adjunctive treatment for symptom control should be considered (Qin 2016 - SR of 7 RCTs: 3 high quality studies, 1 moderate and 3 low)(97)
<b>Constipation</b>	Not reviewed	<b>Potential positive effect</b>	No updates
<b>Obesity</b>	Not reviewed	<b>Potential positive effect</b>	Acupuncture superior to medication, sham and lifestyle modification; low to very low quality evidence (Esteghamati 2015 - Critical review of 3 SRs)(98)
<b>Restless leg syndrome</b>	Insufficient evidence	<b>Potential positive effect</b>	Insufficient evidence (Bega 2016 – Overview of alternative treatment of restless leg syndrome)(99)
<b>Induction of labour</b>	<b>Insufficient evidence</b>	Not reviewed	Insufficient evidence; included studies ranged from high to low quality evidence (Smith 2013 - Cochrane update with 14 RCTs)(100) Acupressure may reduce the duration of labour especially the first stage; insufficient evidence (Mollart 2015 – SR of 7 RCTs)(101)
<b>Bell's palsy</b>	<b>Insufficient evidence</b>	Not reviewed	Seems to be effective; insufficient evidence of efficacy and safety; low quality evidence (Li 2015 – SR & MA of 14 RCTs)(102) Acupuncture superior to waitlist in physical and social function (Kwon 2015 - RCT acupuncture (n=36) vs waitlist (n=13))(103)
<b>Glaucoma</b>	<b>Insufficient evidence</b>	Not reviewed	Insufficient evidence; low quality evidence (Law 2013 - Cochrane SR; 1 RCT on auricular acupressure)(104)
<b>Uterine fibroids</b>	<b>Insufficient evidence</b>	Not reviewed	Acupuncture may be a treatment option as part of a CAM treatment approach (Dalton-Brewer 2016 - Narrative review of CAM)(105)
<b>Vascular dementia</b>	<b>Insufficient evidence</b>	Not reviewed	Acupuncture plus other therapies significantly improved Mini Mental State scores; low quality evidence (Cao 2013 – SR of 12 RCTs)(106)
<b>Erectile dysfunction</b>	Not reviewed	<b>Unclear</b>	Insufficient evidence; low quality evidence (Cui 2016 - SR of 3 RCTs)(107)
<b>Tinnitus</b>	Not reviewed	<b>Unclear</b>	EA - Insufficient evidence; low quality evidence (He 2016 - SR of 5 RCTs)(108)

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which remain unchanged	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
<b>Opiate addiction</b>	Not reviewed	<b>Unclear</b>	Acupuncture is effective in opiate detoxification (and associated depression and anxiety) which is mediated by endogenous dynorphin; no effect on opioid cravings; low quality evidence (Wu 2016 – Narrative review)(109) Acupuncture superior to controls for withdrawal/craving and anxiety post-treatment but not long term; low quality evidence (Grant 2016 - SR & MA of 41 RCTs on substance use disorders)(110)
<b>Chronic fatigue syndrome</b>	Not reviewed	<b>Unclear</b>	Acupuncture plus usual care may improve fatigue in chronic fatigue syndrome and idiopathic chronic fatigue (Kim 2015 - Multi-centre nonblinded RCT, n = 150)(111)
<b>Fibromyalgia</b>	Not reviewed	<b>Unclear</b>	Acupuncture superior to no treatment or standard care in reducing pain and stiffness; low to moderate quality evidence Acupuncture not different from sham in reducing pain, fatigue or improving sleep and global wellbeing; moderate quality evidence EA is probably better than manual acupuncture in reducing pain and stiffness and improving global well-being, sleep and fatigue. (Deare 2013 – Cochrane SR of 9 RCTs)(112) Insufficient evidence; low quality evidence (Yang 2014 – SR of RCTs)(113) Inconsistent evidence (Lauche 2015 – SR of reviews; 2 high and 2 low quality reviews)(114)
<b>Premenstrual syndrome</b>	Not reviewed	<b>Unclear</b>	A SR in 2014 included no new evidence (115) Insufficient evidence (Hofmeister 2016 - Narrative review) (116)
<b>Assisted conception in ART</b>	<b>Insufficient evidence</b> when used around embryo transfer; <b>Not effective</b> when used around oocyte retrieval	Not reviewed	No evidence that acupuncture improves live birth or pregnancy rates in ART regardless of whether performed around the time of oocyte retrieval or embryo transfer (Cheong 2013 - Cochrane update with 20 RCTs)(117). Acupuncture ineffective when used only on the day of oocyte retrieval but effective when used on the day of embryo transfer (Shen 2015 - SR & MA) (118). Acupuncture improves clinical pregnancy rates in women undergoing IVF. Optimal positive effects were seen when acupuncture was used during controlled ovarian hyperstimulation. (Qian 2016 – SR of 30 RCTs & MA)(119) Acupuncture may increase the clinical pregnancy rate and ongoing pregnancy rate and decrease the risk of ovarian hyperstimulation syndrome in women with PCOS undergoing IVF or intracytoplasmic sperm injection (Jo 2017 – SR of 4 RCTs & MA)(120)

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which remain unchanged	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
Improving exercise performance and post-exercise recovery	Not reviewed	<b>Unclear</b>	No updates
Rheumatoid arthritis	Insufficient evidence	<b>Unclear</b>	A SR in 2016 included no new evidence (Fernandez-Llanio Camella 2016 -SR of CAM)(121)
Nausea in pregnancy	Not reviewed	<b>Evidence of no effect</b>	No significant difference between real and sham acupuncture; low quality evidence (Matthews 2015 - Cochrane SR of interventions for nausea in pregnancy; 2 RCTs on acupuncture)(122)
Alcohol dependence	Not reviewed	<b>Evidence of no effect</b>	Low to very low quality evidence (Grant 2016 - SR & MA of 41 RCTs: 12 opioids, 9 cocaine, 11 alcohol, 9 mixed/not reported/other )(110)
Cocaine addiction	Insufficient evidence	<b>Evidence of no effect</b>	Low to very low quality evidence (Grant 2016 - SR & MA of 41 RCTs: 12 opioids, 9 cocaine, 11 alcohol, 9 mixed/not reported/other )(110)
Epilepsy	<b>Not effective</b>	Not reviewed	No evidence of effect (Cheuk 2014 – Cochrane update)(123)

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which have increased in evidence level	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
<b>Low back pain</b>	Possibly effective	Unclear	<b><i>Research has shown better outcomes for chronic low back pain than for acute low back pain</i></b>
Chronic low back pain			High quality evidence for acupuncture; moderate quality evidence for acupuncture (Wellington 2014 – SR of noninvasive interventions)(48) Acupuncture alone or as an adjunct to usual care provided short-term improvement in pain and function; low to high quality evidence; ‘should be advocated in routine clinical practice’ (Liu 2015 - Overview of 16 SRs)(49) Moderate quality evidence for pain and function (Chou 2016 [AHRQ Comparative Effectiveness review])(45) <b><i>Upgrade to positive</i></b>
Acute low back pain			Acupuncture superior to NSAIDs for improving symptoms; acupuncture superior to sham for pain but not function (Lee 2013 - SR of 11 RCTs)(47) Low quality evidence for pain and function (Chou 2016 - AHRQ Comparative Effectiveness review)(45) <b><i>Upgrade to potential positive</i></b>
<b>Headache</b> (frequent episodic or chronic tension-type)	Not reviewed	<b>Potential positive effect</b>	Acupuncture is cost effective for headache (Kim 2012)(76) A potentially important role for acupuncture as part of a treatment plan for migraine, tension-type headache, and several different types of chronic headache disorders. Cost-effective in Germany and UK (Coytaux 2016 - Brief review of selected SRs and MAs)(44) Acupuncture can reduce workplace headache pain intensity, frequency and related disability; low quality evidence (Lardon 2016 - SR of 15 RCTs)(124) Effective for frequent episodic or chronic tension-type headaches; Moderate or low quality evidence (Linde 2016 - Cochrane SR of 12 RCTs)(43) Acupuncture has been included in the NICE guidelines for headache since 2012(38) <b><i>Upgrade to positive</i></b>
<b>Knee osteoarthritis</b>	Not reviewed	<b>Potential positive effect</b>	Acupuncture was equal to balneotherapy and superior to sham acupuncture, muscle-strengthening exercise, Tai Chi, weight loss, standard care and aerobic exercise (in ranked order). Acupuncture superior to standard care and muscle-strengthening exercises in sub-analysis of moderate to high quality studies (Corbett 2013: Network meta-analysis (152 studies on 22 interventions: 12 RCTs included in network MA) (52) <b><i>Upgrade to positive</i></b>

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which have increased in evidence level	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
<b>Postoperative nausea and vomiting</b>	Insufficient evidence	<b>Potential positive effect</b>	Acupuncture and acupressure reduced number of cases of early nausea and vomiting (up to 24 hours); low quality evidence (Cheong 2013 - SR of 30 RCTs)(56) PC 6 stimulation was superior to sham; low quality evidence; no difference between PC 6 stimulation and anti-emetics (moderate quality evidence); insufficient evidence that PC 6 plus anti-emetics is superior to anti-emetics alone (Lee 2015 - Cochrane update with 59 RCTs of PC 6 stimulation)(16) <b>Upgrade to positive</b>
Postoperative nausea and vomiting and pain after tonsillectomy			Pain scores, analgesic need and nausea and vomiting were reduced (acupuncture vs control); no significant adverse events; insufficient evidence (Cho 2016 - MA of 12 RCTs; 7 high quality)(125) Acupuncture superior to controls and cost-effective; insufficient evidence (Shin 2016 - SR & MA - 4 RCTs [3 high quality]; 3 randomised prospective studies and 1 pilot)(126) <b>Upgrade to positive</b>
<b>Postoperative pain</b>	Not reviewed	<b>Unclear</b>	Some forms of acupuncture (acupuncture, EA and TEAS) improved pain on day 1 after surgery and reduced opioid use; subgroup analysis showed acupuncture and TEAS superior to EA; moderate quality evidence (Wu 2016 - SR & MA of 13 RCTs)(17) <b>Upgrade to positive</b>
Postoperative pain – back surgery			Encouraging but limited evidence (Cho 2015 - SR of 5 RCTs; 3 high quality)(57) <b>Upgrade to positive</b>
Postoperative pain – knee surgery			Acupressure reduced pain; acupuncture did not reduce pain but resulted in reduced use of ibuprofen; low quality evidence (Barlow 2013 - SR of CAM – 3 acupuncture & 1 acupressure RCTs)(58) Acupuncture superior to sham in post-operative fentanyl use, time to first request for fentanyl and pain intensity (Chen 2015 - RCT acupuncture vs sham, n=60)(59) <b>Upgrade to positive</b>

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which have increased in evidence level	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
<b>Stroke</b>	<b>Insufficient evidence</b>	Not reviewed	<b><i>Stroke research has now become nuanced into several new topics</i></b> Acupuncture may be effective for treating post-stroke neurological impairment and dysfunction such as dysphagia, but does not prevent post-stroke death; (Zhang 2014 - Review of SRs: 8 reviews high quality, 6 moderate, 10 low)(127) <b><i>Upgrade to potential positive</i></b>
Acute stroke			Acupuncture plus rehabilitation superior to rehabilitation alone for acute and subacute stroke sequelae (Vados 2015 – SR of 17 RCTs; 5 high quality)(128) Acute ischaemic stroke: EA superior to usual care in Barthel Index, Fugl-Meyer Assessment, National Institutes of Health Stroke Scale, and Revised Scandinavian Stroke Scale; Moderate to high quality evidence (Liu 2015 - MA of 18 RCTs)(129) <b><i>Upgrade to potential positive</i></b>
Stroke rehabilitation			May have beneficial effects on improving dependency, global neurological deficiency, and some specific neurological impairments; insufficient evidence; low to very low quality evidence (Yang 2016 – Cochrane update)(130) <b><i>Upgrade to potential positive</i></b>
Post-stroke spasticity			Acupuncture or EA significantly decreased spasticity after stroke; low quality evidence (Lim 2015 – SR & MA of 5 RCTs)(131) Acupuncture improved passive resistance to stretching, degree of personal dependence and motor function; insufficient evidence; 6 high quality studies; 3 moderate quality (Rodriguez-Mansilla 2016 – SR of 9 RCTs or controlled studies)(132) <b><i>Upgrade to potential positive</i></b>
Post-stroke insomnia			Acupuncture superior to sham acupuncture and to medication; insufficient evidence; low to moderate evidence (Lee 2016 – SR & MA of 13 RCTs)(133) <b><i>Upgrade to potential positive</i></b>
Post-stroke shoulder pain			Acupuncture plus rehabilitation superior to rehabilitation alone; insufficient evidence; low to moderate evidence (Lee 2016 – SR of 12 RCTs)(134) <b><i>Upgrade to potential positive</i></b>
Post-stroke hiccoughs			Acupuncture may be effective as an adjunctive but not as a stand-alone treatment; low quality evidence (Yue 2016 – SR & MA of 5 RCTs)(135) <b><i>Unclear</i></b>

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which have increased in evidence level	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
<b>Menopausal symptoms</b>	Not reviewed	<b>Unclear</b>	<b><i>Menopausal symptom research has now separated into two new topics</i></b>
Menopausal hot flushes			Acupuncture improves hot flash frequency and severity, menopause-related symptoms, and QoL (vasomotor domain) in natural menopause (Chiu 2015 - MA of 12 studies; 2 high quality studies, 8 moderate and 2 low)(136) Promising results; low quality evidence (Selva Ovid 2013 (Review of 8 SRs and 9 RCTs))(93) Acupuncture superior to wait-list or no treatment; low quality evidence Acupuncture superior to sham in reducing severity but not frequency; very low quality evidence Acupuncture inferior to hormone therapy in QoL and frequency; no difference in severity; low quality evidence (Dodin 2013 - Cochrane SR of 16 RCTs)(137) <b><i>Upgrade to potential positive</i></b>
Perimenopausal and postmenopausal sleep disturbance			Significant reduction in sleep disturbance which appears to be associated with changes in serum estradiol, FSH and LH; acupuncture recommended as adjunctive therapy in improving sleep disturbances in perimenopausal and postmenopausal women (Chiu 2016 - SR of 34 studies; 4 high quality)(138) Improved sleep quality; limited evidence; moderate to high quality evidence (Bezerra 2015 - SR of 7 RCTs; 4 high quality; no studies with high risk of bias)(139) <b><i>Upgrade to potential positive</i></b>
<b>Cancer adverse effects</b> (not including cancer pain and chemotherapy-induced nausea and vomiting)	Not reviewed	<b>Unclear</b>	<b><i>Cancer adverse effects research has now become nuanced into several new topics</i></b> Acupuncture plus usual care superior to usual care alone in reducing pain, fatigue, and in improving QoL. Acupuncture's effectiveness for managing anorexia, reducing constipation, paraesthesia and dysaesthesia, insomnia, and limb oedema is unclear; low to very low quality evidence (Lau 2016 - SR & MA of 13 RCTs)(140) Acupuncture may be effective for cancer pain, post-operative pain, aromatase inhibitor related joint pain and neck dissection pain and dysfunction, as well as opioid related constipation and pruritus and chemotherapy-induced nausea and vomiting and neuropathy; no assessment of quality of evidence (Lu 2013 - Narrative review)(141) Acupuncture effective for treatment-related nausea and vomiting, cancer pain, fatigue, hot flushes, xerostomia, dyspnoea and anxiety; low quality evidence (Towler 2013 - Review of 17 reviews)(142) Acupuncture may be appropriate adjunctive therapy for a range of cancer-related symptoms including adverse effects of chemotherapy and radiotherapy and cancer pain (Lian 2014 - SR of 33 RCTs)(143)

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which have increased in evidence level	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
Cancer-related fatigue			<p>Acupuncture and moxibustion appear to be efficacious adjunctive therapy; Insufficient evidence; low quality evidence (He 2013 - SR of 7 RCTs)(144)</p> <p>Acupuncture and acupressure tend to be effective, acupuncture more than acupressure; low quality evidence (Ling 2013 - SR)(145)</p> <p>Conflicting evidence: 4 studies showed acupuncture or acupuncture plus usual care superior to sham, usual care, enhanced usual care or no treatment; 3 studies showed no difference between acupuncture and sham; very low quality evidence (Posadzki 2013 - SR of 7 RCTs)(146)</p> <p>Acupuncture may reduce fatigue after cancer treatment; low quality evidence (Finnegan-John 2013 - SR of CAM 20 studies; 3 acupuncture/acupressure RCTs)(147)</p> <p>Acupuncture plus education superior to usual care; low quality evidence (Zeng 2014 (SR of 7 RCTs)(148)</p> <p><b>Upgrade to potential positive</b></p>
Cancer-related insomnia			<p>Acupuncture may be superior to sham acupuncture, drugs or hormones therapy. Number of studies and effect size are small for clinical significance; low quality evidence (Choi 2016 - SR of 6 RCTs)(149)</p> <p><b>Unclear</b></p>
Hot flushes/flushes in breast cancer			<p>Acupuncture superior to sham in some studies and superior to baseline in all studies; low quality evidence (Garcia 2015 - SR of 8 RCTs)(150)</p> <p>Acupuncture had similar efficacy to venlafaxine and gabapentin but may have longer durability after completing treatment and fewer side effects (Johns 2016 - SR of interventions; 2 acupuncture vs medication studies)(151)</p> <p>Acupuncture superior to sham in 3 studies; no different from sham in 6 studies; inferior to hormone therapy in 2 studies; low quality evidence (Chen 2016 - SR of 12 RCTs)(152)</p> <p>Conflicting evidence; low quality evidence (Salehi 2016 - SR of 12 studies)(153)</p> <p>Acupuncture yielded small-size effects on reducing hot-flash frequency and the severity of menopause-related symptoms; Insufficient or conflicting evidence; Low to high quality studies (Chiu 2016 - SR of 7 studies; 4 high quality studies)(87)</p> <p><b>Unclear</b></p>

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which have increased in evidence level	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
Xerostomia in cancer	Not reviewed	<b>Unclear</b>	Insufficient evidence (Zhuang 2013 - SR of 4 studies)(154) Small increase in saliva production; low quality evidence (Furness 2013 - Cochrane SR of non-pharmacological interventions - 9 RCTs; 5 acupuncture)(155) Acupuncture was superior to sham; low quality evidence (Hanchanale 2015 - SR of 6 RCTs)(156) <b>Unclear</b>
Recovery after colorectal cancer resection			Acupuncture efficacious and effective; Low to moderate quality evidence (Kim 2016 - SR of 7 RCTs)(157) <b>Upgrade to potential positive</b>
Aromatase-inhibitor-induced arthralgia			Acupuncture superior to sham in 2 high quality studies; no different from sham in 2 low-quality studies (Bae 2015 - SR of 4 RCTs)(158) Acupuncture reduces joint pain and stiffness but not superior to sham; moderate to high quality evidence (Chien 2015 - SR of 5 RCTs)(159) <b>Upgrade to potential positive</b>
Chemotherapy-induced peripheral neuropathy			Acupuncture superior to sham in one RCT; very low quality evidence (Franconi 2013 - SR of 3 RCTs, 3 case series, 1 rat study)(160) <b>Unclear</b>
Myelosuppression after chemotherapy			Insufficient evidence; low to very low quality evidence (Fu 2015 - Narrative review includes 7 RCTs)(161) <b>Unclear</b>
Cancer-related psychological symptoms			All included studies suggest benefits in depression, anxiety, sleep disturbance, and for improving QoL; strong evidence for safety; no assessment of quality of evidence (Haddad 2014 - SR of 12 studies; 8 RCTs)(162) <b>Unclear</b>
<b>Surgical analgesia</b>	Not reviewed	<b>Unclear</b>	<b><i>Surgical analgesia has now become nuanced into several new topics</i></b> Conflicting evidence; insufficient evidence; no assessment of quality of evidence (Lee 2014 - Overview of 12 SRs on postsurgical nausea and vomiting and postsurgical pain)(163)

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which have increased in evidence level	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
Craniotomy anaesthesia			Acupuncture significantly reduced the amount of volatile anaesthetics and led to faster extubation time and postoperative patient recovery and reduced postoperative nausea and vomiting. In addition, significantly reduced blood levels of the brain tissue injury marker S100beta 48 hours after operation; low quality studies excluded from MA, hence moderate to high quality evidence (Asmussen 2016 - MA of 10 RCTs)(164) <b>Upgrade to potential positive</b>
Ambulatory anaesthesia			Acupuncture may reduce preoperative anxiety, and postoperative pain, nausea, vomiting, shivering and emergence delirium. Acupuncture is safe and cost-effective. Acupuncture may be a beneficial adjunctive therapy for ambulatory anaesthesia; insufficient evidence; high quality studies favour acupuncture (Liodden 2013 - Narrative review)(165) <b>Upgrade to potential positive</b>
Postoperative nausea and vomiting and pain after tonsillectomy			Pain scores, analgesic need and nausea and vomiting were reduced (acupuncture vs control); no significant adverse events; insufficient evidence (Cho 2016 - MA of 12 RCTs; 7 high quality)(125) Acupuncture superior to controls and cost-effective; insufficient evidence (Shin 2016 - SR & MA - 4 RCTs [3 high quality]; 3 randomised prospective studies and 1 pilot)(126) <b>Upgrade to potential positive</b>
Dry eye	Not reviewed	<b>Unclear</b>	Acupuncture was significantly superior to artificial tears in tear break-up time, Schirmer I test, and cornea fluorescein staining; Low to moderate quality evidence (Yang 2015 - SR & MA of 7 RCTs)(166) <b>Upgrade to potential positive</b>

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which have increased in evidence level	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
<b>Carpal tunnel syndrome</b>	Not reviewed	<b>Evidence of no effect</b>	Acupuncture superior to ibuprofen in function and symptoms (Hadianfard et al 2015 - RCT comparing acupuncture plus night wrist splints with ibuprofen plus night wrist splints, n=50)(167) <b>Upgrade to unclear</b>
<b>Irritable bowel syndrome</b>	Insufficient evidence	<b>Unclear</b>	Insufficient evidence; low to moderate quality evidence (Manheimer 2012 - Cochrane SR of 17 RCTs)(168) Acupuncture superior to usual care on IBS symptom severity score at 6, 9 and 12 months but not at 24 months (MacPherson 2016 - high quality RCT)(169) <b>Upgrade to potential positive</b>
<b>Hypertension</b>	Not reviewed	<b>Unclear</b>	Acupuncture plus medication superior to sham plus medication; low quality evidence (Wang 2013 - SR of 35 RCTs)(170) Acupuncture plus medication superior to medication, but acupuncture not superior to medication; high quality evidence (Li 2014 - SR of 4 RCTs) (171) Acupuncture plus medication superior to medication, but acupuncture not superior to medication; risk of bias unclear for most domains (Zhao 2015 – 23 RCTs)(172) <b>Upgrade to potential positive</b>
<b>Lateral elbow pain</b>	<b>Insufficient evidence</b>	Not reviewed	Acupuncture superior to sham; moderate quality evidence (Gadau 2014 - SR of 19 RCTs)(173) Insufficient evidence: low to very low quality evidence (Tang 2015 - SR of 4 RCTs)(174) <b>Upgrade to potential positive</b>
<b>Neck pain</b>	Insufficient evidence	<b>Unclear</b>	Acupuncture plus usual medical care is cost-effective for neck pain and its associated disorders (NAD) (Van der Velde 2015 – SR of 6 studies)(175) Acupuncture superior to sham acupuncture or inactive treatment (at completion of treatment and short-term follow-up) for pain relief; Moderate quality evidence (Trinh 2016 - Cochrane update with 27 RCTs)(176) Limited evidence of effectiveness; low quality evidence (Moon 2014 – 6 RCTs on whiplash associated disorder [WAD])(177) <b>Moderate quality evidence for NAD, but low quality evidence for WAD</b> <b>Upgrade to potential positive for NAD, but leave WAD as unclear</b>

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which have increased in evidence level	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
<b>Shoulder pain</b>	Insufficient evidence	<b>Unclear</b>	For non-operative treatment options at an early stage of Shoulder Impingement Syndrome (SIS), exercise combined with therapies such as kinesio taping, specific exercises, and acupuncture should be considered as the first line choices (2 included high quality acupuncture studies) (Dong 2015 - SR and network MA)(178)  <i>Upgrade to potential positive</i>
<b>Asthma</b>	<b>Insufficient evidence</b>	Not reviewed	Acupuncture may improve peak expiratory flow or peak expiratory flow variability in children; low quality evidence (Liu 2015 – SR of 7 RCTs)(179) Acupoint herbal patches superior to sham in improving FEV1 and asthma symptoms; low quality evidence (Lee 2016 – SR & MA of 16 RCTs)(180) Acupuncture improved FEV1, FEV1/FVC, IL and IgE (moderate quality evidence), but not ECP (Su 2016 - SR & MA of 8 RCTs)(181)  <i>Upgrade to potential positive</i>

**Table 5. Evidence levels from 2005 to 2016: explanatory notes with references (contd)**

Conditions which have decreased in evidence level	Australian DVA (Sept 2005 - Sept 2010)	USVA (Jan 2005 - Mar 2013)	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)
	<i>Baseline evidence in bold</i>		
Dysmenorrhoea	Not reviewed	<b>Potential positive effect</b>	<p><b>Conflicting SRs (3 positive SRs; Cochrane SR insufficient evidence)</b></p> <p>Acupuncture is effective and acupressure may be effective for pain relief; acupuncture trials had low to moderate risk of bias; acupressure trials high risk of bias (Chen et al, 2013 – MA of 3 acupuncture and 4 acupressure RCTs)(182)</p> <p>Acupoint stimulation superior to controls for pain relief; low to moderate quality evidence (Xu et al, 2014 MA of 20 RCTs of acupoint stimulation)(183)</p> <p>Acupuncture and acupressure (vs placebo, waitlist or medication) reduced pain intensity, while acupuncture also improved quality of life; moderate quality evidence (Abaraogu 2015 – SR of 8 RCTs and MA of 4 RCTs)(184)</p> <p>Insufficient evidence; low to very low quality evidence (Smith 2016 - Cochrane update)(185)</p> <p><b><i>Downgrade to unclear</i></b></p>
Ankle sprain	Not reviewed	<b>Potential positive effect</b>	<p>Insufficient evidence; low to very low quality evidence (Kim 2014 – Cochrane SR of 15 RCTs, 5 quasi-randomised trials)(186)</p> <p><b><i>Downgrade to unclear</i></b></p>

**Table 6. All conditions reviewed sorted by clinical areas**

Condition	<p style="text-align: center;"><b>THE ACUPUNCTURE EVIDENCE PROJECT</b>  <b>(updates from March 2013 to August 2016)</b>  <b>systematic reviews, meta-analyses, network meta-analyses,</b>  <b>narrative reviews, cost effectiveness reviews</b>  <i><b>Cochrane reviews highlighted in bold</b></i></p>	Comments
<b>Cardiovascular/respiratory conditions</b>		
Angina pectoris	<p>Xu 2013 (Narrative review)(187): Acupuncture shows effectiveness rates of 80% to 92.6% without adverse effects of medication; no assessment of quality of evidence</p> <p>Yu 2015 (SR of 25 RCTs)(188): Improvement in symptoms and ECG; very low quality evidence</p> <p>Zhang 2015 (SR &amp; MA of 15 RCTs)(189): Acupuncture plus medication superior to medication alone in improving symptoms and ECG; insufficient evidence; no assessment of quality of evidence</p>	<i><b>Unclear</b></i>
Asthma	<p>Liu 2015 [Childhood asthma] (SR of 7 RCTs)(179): Acupuncture may improve peak expiratory flow or peak expiratory flow variability; insufficient evidence; low quality evidence</p> <p>Lee 2016 [Acupoint herbal patches] (SR &amp; MA of 16 RCTs)(180): Superior to sham in improving FEV1 and asthma symptoms; low quality evidence</p> <p>Su 2016 [Acupoint application in adults] (SR &amp; MA of 8 RCTs)(181): Acupuncture improved FEV1, FEV1/FVC, IL and IgE (moderate quality evidence) but not ECP</p>	<p>Moderate quality evidence of lung function improvements in adults -<i><b>Potential positive</b></i> (adults)</p> <p><i><b>Unclear</b></i> (children)</p>
Chronic obstructive pulmonary disease (COPD)	Coyle 2014 (SR of 16 RCTs)(190): Clinically significant improvements in QoL and dyspnoea, but not lung function; moderate quality evidence	<i><b>Unclear</b></i>
Heart failure	Lee 2016 (SR of 7 RCTs)(191): Acupuncture improved exercise capacity, quality of life, hemodynamic parameters, and time domain heart rate variability parameters; low quality evidence	<i><b>Unclear</b></i>
Hypertension	<p>Wang 2013 (SR of 35 RCTs)(170): Acupuncture plus medication superior to sham plus medication; low quality evidence</p> <p>Li 2014 (SR of 4 RCTs)(171): Acupuncture plus medication superior to medication, but acupuncture not superior to medication; high quality evidence</p> <p>Zhao 2015 (SR &amp; MA of 23 RCTs)(172): Acupuncture plus medication superior to medication, but acupuncture not superior to medication; risk of bias unclear for most domains</p>	<p>Low to high quality evidence for acupuncture plus medication</p> <p><i><b>Potential positive</b></i> (with medication)</p>
Vascular cognitive impairment without dementia	Min 2016 (MA of 15 studies)(192): Acupuncture superior to usual care or medication; low quality evidence	<i><b>Unclear</b></i>

<b>Table 6. All conditions reviewed sorted by clinical areas (contd)</b>		
<b>Condition</b>	<b>THE ACUPUNCTURE EVIDENCE PROJECT</b>	<b>Comments</b>
<b>Cardiovascular/respiratory conditions (contd)</b>		
Vascular dementia	Cao 2013 (SR of 12 RCTs)(106): Acupuncture plus other therapies significantly improved Mini Mental State scores; low quality evidence	<i>Unclear</i>
<b>Musculoskeletal disorders</b>		
Acute ankle sprain in adults	Park 2013 (SR of 17 RCTs; 2 high quality, 15 low quality)(193): Acupuncture superior to various controls in relieving pain, facilitating return to normal activity, and promoting QoL based on subgroup analysis of 2 high quality studies <b>Kim 2014 (Cochrane SR of 15 RCTs, 5 quasi-randomised trials)(186):</b> Insufficient evidence; Low to very low quality evidence	<b>Potential positive</b> in USVA review 2014; Insufficient evidence/ low to very low quality evidence (Cochrane) <b>Downgrade to unclear</b>
Carpal tunnel syndrome	Hadianfard 2015 (RCT comparing acupuncture plus night wrist splints with ibuprofen plus night wrist splints, n=50)(167): Acupuncture superior to ibuprofen in function and symptoms	USVA review 2014 ‘Evidence of no effect’; 1 positive RCT in 2015 <b>Upgrade to unclear</b>
Fibromyalgia	<b>Deare 2013 (Cochrane SR of 9 RCTs)(112):</b> Acupuncture superior to no treatment or standard care in reducing pain and stiffness; low to moderate quality evidence Acupuncture not different from sham in reducing pain, fatigue or improving sleep and global wellbeing; moderate quality evidence EA is probably better than manual acupuncture in reducing pain and stiffness and improving global well-being, sleep and fatigue. Yang 2014 (MA of 9 RCTs)(113): Insufficient evidence; low quality evidence Lauche 2015 (SR of reviews; 2 high and 2 low quality reviews)(114): Inconsistent evidence	<i>Unclear</i>
Lateral elbow pain	Gadau 2014 (SR of 19 RCTs)(173): Acupuncture superior to sham; moderate quality evidence Tang 2015 (SR of 4 RCTs)(174): Insufficient evidence: low to very low quality evidence	Moderate quality evidence when 19 RCTs included – <b>Upgrade to potential positive</b>
Lumbar spinal stenosis	Kim 2013 (SR of 6 RCTs, 6 controlled trials) (194): Acupuncture superior to controls in pain intensity, functional improvements and QoL; Low quality evidence	<i>Unclear</i>

**Table 6. All conditions reviewed sorted by clinical areas (contd)**

Condition	THE ACUPUNCTURE EVIDENCE PROJECT	Comments
<b>Musculoskeletal disorders (contd)</b>		
Low back pain	Taylor 2014 (51): Cost effective for chronic low back pain Andronis 2016 (50): Likely to be cost effective	Chronic and acute low back pain need to be differentiated in SRs
Acute low back pain	Lee 2013 (SR of 11 RCTs; 5 low risk of bias, 6 high)(47): Acupuncture superior to NSAIDs for improving symptoms; acupuncture superior to sham for pain but not function; Chou 2016 [AHRQ Comparative Effectiveness review])(45): Low quality evidence for pain and function	<b>Upgrade to potential positive</b>
Chronic low back pain	Lam 2013 (32 RCTs; 25 in MA)(46): Acupuncture may be effective for pain and functional limitation in chronic non-specific low back pain: Evidence Level 1 Wellington 2014 (SR of non-invasive interventions)(48): High quality evidence for acupuncture; moderate quality evidence for acupuncture Liu 2015 (Overview of 16 SRs)(49): Acupuncture alone or as an adjunct to usual care provided short-term improvement in pain and function; low to high quality evidence; 'should be advocated in routine clinical practice' Chou 2016 [AHRQ Comparative Effectiveness review])(45): Moderate quality evidence for pain and function Nahin 2016 (4 RCTs; Excluded studies not performed in USA or by US researchers)(195): Acupuncture superior to usual care; Acupuncture superior to sham in 1 RCT, but not superior in 2 RCTs	Moderate to high quality evidence Cost effective, safe <b>Upgrade to positive</b>
Neck pain	Van der Velde 2015 (SR of 6 studies)(175): Acupuncture plus usual medical care is cost-effective for neck pain and its associated disorders (NAD) <b>Trinh 2016 (Cochrane update with 27 RCTS)(176):</b> Acupuncture superior to sham acupuncture or inactive treatment (at completion of treatment and short-term follow-up) for pain relief; Moderate quality evidence	Moderate quality evidence (Cochrane update); Acupuncture plus medication is cost-effective <b>Potential positive for NAD</b>
Knee osteoarthritis pain	Corbett 2013 (Network meta-analysis - 152 studies on 22 interventions: 12 RCTs included in network MA)(52): Acupuncture was equal to balneotherapy and superior to sham acupuncture, muscle-strengthening exercise, Tai Chi, weight loss, standard care and aerobic exercise (in ranked order). Acupuncture superior to standard care and muscle-strengthening exercises in sub-analysis of moderate to high quality studies Nahin 2016 (4 RCTs; Excluded studies not performed in USA or by US researchers)(195): Acupuncture superior to attention control or usual care in 3/4 studies; Acupuncture superior to sham in 2/4 studies	Moderate to high quality evidence <b>Upgrade to positive</b>

<b>Table 6. All conditions reviewed sorted by clinical areas (contd)</b>		
<b>Condition</b>	<b>THE ACUPUNCTURE EVIDENCE PROJECT</b>	<b>Comments</b>
<b>Musculoskeletal disorders (contd)</b>		
Osteoarthritis	Kim 2012 (Cost effectiveness analysis)(76): Acupuncture is cost effective for dysmenorrhoea, allergic rhinitis, osteoarthritis & headache Manyanga 2014 (SR & MA of 12 trials)(53): Significant reductions in pain intensity, improvement in functional mobility and quality of life; low quality evidence	Cost-effective
Plantar heel pain		<b>Potential positive</b> in USVA review 2014; no updates
Restless leg syndrome	Bega 2016 (Overview of alternative treatment of restless leg syndrome)(99): insufficient evidence	<b>Potential positive</b>
Sciatica	Lewis 2015 (Network MA of 21 interventions)(196): Acupuncture 2 <sup>nd</sup> out of 21 interventions for global effect and pain intensity Qin 2015 (SR & MA of 11 RCTs; 10 acupuncture vs medications; 1 acupuncture vs sham)(197): Acupuncture may be superior to drugs and may enhance the effect of drugs for patients with sciatica; low quality evidence Ji 2015 (SR of 12 RCTs)(198): Acupuncture superior to conventional Western medicine in outcomes effectiveness, pain intensity and pain threshold; low quality evidence	Acupuncture 2 <sup>nd</sup> out of 21 interventions for global effect and pain intensity <b>Potential positive</b>
Shoulder pain: Shoulder impingement syndrome (SIS)	Dong 2015 (SR and network MA; 2 high quality acupuncture studies)(178): For non-operative treatment options at an early stage of SIS, exercise combined with therapies such as kinesio taping, specific exercises, and acupuncture should be considered as the first-line choices, whereas pulsed electromagnetic field therapy, localized corticosteroid injection, diacutaneous fibrolysis, and ultrasound therapy may be considered as the second-line treatment choices; however, low-level laser therapy and the localized injection of NSAIDs are not recommended	Moderate to high quality evidence <b>Potential positive</b>
Whiplash associated disorder (WAD)	Moon 2014 (SR of 6 RCTs)(177): Limited evidence of effectiveness; low quality evidence	<b>Unclear for WAD</b>

**Table 6. All conditions reviewed sorted by clinical areas (contd)**

Condition	THE ACUPUNCTURE EVIDENCE PROJECT	Comments
<b>Neurological disorders</b>		
Acute stroke	Vados 2015 (SR of 17 RCTs; 5 high quality)(128) Acupuncture plus rehabilitation superior to rehabilitation alone for acute and subacute stroke sequelae Liu 2015 (MA of 18 RCTs)(129): Acute ischaemic stroke: EA superior to usual care in Barthel Index, Fugl-Meyer Assessment, National Institutes of Health Stroke Scale, and Revised Scandinavian Stroke Scale; Moderate to high quality evidence Liu 2015: (Acute ischaemic stroke) EA superior to usual care in Barthel Index, Fugl-Meyer Assessment, National Institutes of Health Stroke Scale, and Revised Scandinavian Stroke Scale; Moderate to high quality evidence	Moderate to high quality evidence <b>Potential positive</b>
Bell's palsy	Li 2015 (SR & MA of 14 RCTs)(102): Seems to be effective; insufficient evidence of efficacy and safety; low quality evidence Kwon 2015 (RCT acupuncture (n=36) vs waitlist (n=13))(103): RCT acupuncture vs waitlist (n=26 vs 13) Acupuncture superior to waitlist in physical and social function	<b>Unclear</b>
Epilepsy	<b>Cheuk 2014 (Cochrane update)(123)</b> : No evidence of effect; low quality evidence	<b>No evidence of effect</b> (Cochrane update)
Parkinson's disease	Kim 2014 (Review of 11 studies: 6 RCTs, 4 uncontrolled open label studies & 1 crossover trial)(199): Insufficient evidence; low to very low quality evidence	<b>Unclear</b>
Post-stroke spasticity	Lim 2015 (SR & MA of 5 RCTs)(131): Acupuncture or EA significantly decreased spasticity after stroke; low quality evidence Rodriguez-Mansilla 2016 (SR of 9 RCTs or controlled studies: 6 high quality; 3 moderate) (132) Acupuncture improved passive resistance to stretching, degree of personal dependence and motor function; insufficient evidence	<b>Potential positive</b>
Stroke	Zhang 2014 (Review of SRs: 8 reviews high quality, 6 moderate, 10 low)(127): Acupuncture may be effective for treating post-stroke neurological impairment and dysfunction such as dysphagia, but does not prevent post-stroke death	Moderate to high quality evidence
Stroke rehabilitation	<b>Yang 2016 (Cochrane update)(130)</b> : May have beneficial effects on improving dependency, global neurological deficiency, and some specific neurological impairments; low to very low quality evidence	<b>Potential positive</b> (Cochrane update)

**Table 6. All conditions reviewed sorted by clinical areas (contd)**

Condition	THE ACUPUNCTURE EVIDENCE PROJECT	Comments
<b>Mental health</b>		
Anxiety	Bazzan 2014 (Narrative review)(79): Acupuncture has ‘potential use’ Goyata 2016 (Integrative review of 19 studies including 6 RCTs; 11 high quality studies; 5 moderate)(80): Positive and statistically significant effects; Moderate to high quality evidence	Moderate to high quality evidence <b>Potential positive</b>
Alzheimers disease	Zhou 2015 (SR & MA of 10 RCTs)(200): Acupuncture superior to medication in improving cognitive function on MMSE. Acupuncture plus medication superior to medication alone. Acupuncture is safe; no assessment of quality of evidence	<b>Unclear</b>
Chronic fatigue syndrome	Kim 2015 (Multi-centre non-blinded RCT, n=150)(111): Acupuncture plus usual care may improve fatigue in chronic fatigue syndrome and idiopathic chronic fatigue	<b>Unclear</b>
Depression	Chan 2015 (SR & MA of 13 RCTs;1 high quality, 5 moderate, 7 low)(88): Acupuncture plus SSRIs superior to SSRIs alone, with an early onset of action and was safe and well-tolerated; EA had greater effect than manual acupuncture Bosch 2015 (Review of SRs & MAs)(82): Effective and safe for major depressive disorder, especially in improving sleep, mood and QoL by modulating and normalizing the limbic–paralimbic–neocortical network (LPNN), including the default mode network (DMN); ‘promising’ evidence Spackman 2014 (Cost-effectiveness analysis)(89): Acupuncture is cost-effective compared with counselling or usual care alone, although the ranking of counselling and acupuncture depends on the relative cost of delivering these interventions	<b>Potential positive</b> in USVA 2014; 2 positive SRs since Cost-effective
Insomnia	Zhao 2013 (SR)(90): Acupuncture may be superior to medication. Acupuncture for insomnia is potentially mediated by norepinephrine, melatonin, gamma-aminobutyric acid, and beta-endorphin; Insufficient evidence Shergis 2016 (SR of 30 studies)(91): Acupuncture statistically superior to sham (3 studies) and medication (27 studies); low quality evidence	<b>Potential positive</b> in USVA 2014; 2 positive SRs since
Opioid detoxification	Wu 2016 (109): Acupuncture is effective in opiate detoxification (and associated depression and anxiety) which is mediated by endogenous dynorphin; no effect on opioid cravings; low quality evidence	<b>Unclear</b>
Post-traumatic stress disorder	Engel 2015 (RCT acupuncture plus usual care vs usual care, n=55)(81): Acupuncture plus usual care superior to usual care in PTSD severity, depression, pain and physical and mental functioning	<b>Potential positive</b> in USVA 2014; positive RCT in 2015
Pre-treatment anxiety	Au 2015 (SR of 7 RCTs, MA of 5)(201): Acupressure effective in reducing anxiety; Moderate to high quality evidence	Moderate to high quality evidence <b>Potential positive</b>

**Table 6. All conditions reviewed sorted by clinical areas (contd)**

Condition	THE ACUPUNCTURE EVIDENCE PROJECT	Comments
<b>Mental health (contd)</b>		
Schizophrenia	<p>Bosch 2015 (Review of SRs &amp; MAs)(82): Acupuncture effective for schizophrenia, especially in improving sleep, mood and QoL by modulating and normalizing the limbic–paralimbic–neocortical network (LPNN), including the default mode network (DMN); limited evidence</p> <p><b>Shen 2014 (Cochrane update of 30 RCTs)(83):</b> Acupuncture plus antipsychotic medication superior to antipsychotic medication alone, in terms of mental state and length of hospitalisation (moderate quality evidence) with fewer adverse effects (low quality evidence)</p>	<p>Moderate quality evidence for acupuncture with antipsychotics</p> <p><b>Potential positive</b></p>
Smoking cessation	<p><b>White 2014 (Cochrane SR Of 38 RCTs)(84):</b> Acupuncture and acupressure superior to sham short-term; low quality evidence</p> <p>Insufficient evidence of effects for 6 months or more</p>	<p><b>Potential positive</b> short term effects</p> <p><b>No evidence of effect</b> after 6 months (Cochrane update)</p>
<b>Gynaecology and obstetrics</b>		
Back or pelvic pain during pregnancy	<p>Selva Olid 2013 (Review of 8 SRs and 9 RCTs)(93): Promising results; low quality evidence</p> <p>Close 2014 (SR of 8 RCTs on CAM; 2 acupuncture RCTs with low risk of bias)(94): Clinically important and statistically significant changes</p> <p><b>Liddle 2015 (Cochrane SR: Comparison of interventions 26 RCTs – 7 acupuncture RCTs)(95):</b> Moderate quality evidence showed that acupuncture or exercise, tailored to the stage of pregnancy, significantly reduced evening pelvic or lumbo-pelvic pain. Acupuncture superior to exercise for reducing evening pelvic pain; Both acupuncture and exercise were superior to usual care; Insufficient evidence</p>	<p><b>Potential positive</b></p>
Dysmenorrhoea	<p>Chen et al, 2013 (MA of 3 acupuncture and 4 acupressure RCTs)(202): Acupuncture is effective and acupressure may be effective for pain relief; acupuncture trials had low to moderate risk of bias; acupressure trials high risk of bias</p> <p>Xu 2014 (MA of 20 RCTs of acupoint stimulation)(183): Acupoint stimulation superior to controls for pain relief; low to moderate quality evidence</p> <p>Abaraogu 2015 (SR of 8 RCTs and MA of 4 RCTs)(184): Acupuncture and acupressure vs placebo, waitlist or medication reduced pain intensity, while acupuncture also improved physical and mental aspects of QoL; moderate quality evidence</p> <p><b>Smith 2016 (Cochrane update)(185):</b> Insufficient evidence; low to very low quality evidence</p> <p>Kim 2012(76): Acupuncture is cost effective for dysmenorrhoea, allergic rhinitis, osteoarthritis &amp; headache</p>	<p><b>Conflicting SRs (3 positive SRs vs Cochrane SR insufficient evidence)</b></p> <p>Moderate quality evidence vs low to very low quality evidence;</p> <p>Cost effective</p> <p><b>Downgrade to unclear</b></p>

<b>Table 6. All conditions reviewed sorted by clinical areas (contd)</b>		
<b>Condition</b>	<b>THE ACUPUNCTURE EVIDENCE PROJECT</b>	<b>Comments</b>
<b>Gynaecology and obstetrics (contd)</b>		
Induction of labour	<b>Smith 2013 (Cochrane SR of 14 RCTs)(100):</b> Insufficient evidence; included studies ranged from high to low quality evidence Mollart 2015 (SR of 7 RCTs)(101): Acupuncture may reduce the duration of labour especially the first stage; insufficient evidence	<b>Unclear</b> (Cochrane update)
Labour pain	Levett 2014 (Critical narrative review of SRs)(92): Acupuncture & acupressure ‘promising’ – Conflicting results due to heterogeneity in study designs, research questions, treatment protocols and outcomes measures	<b>Potential positive</b>
Menopausal hot flushes	Chiu 2015 (MA of 12 studies; 2 high quality, 8 moderate and 2 low)(136): Acupuncture improves hot flash frequency and severity, menopause-related symptoms, and QoL (vasomotor domain) in natural menopause Selva Ovid 2013 (Review of 8 SRs and 9 RCTs) (93): Promising results; low quality evidence <b>Dodin 2013 (Cochrane SR of 16 RCTs)(137):</b> Acupuncture superior to wait-list or no treatment; low quality evidence Acupuncture superior to sham in reducing severity but not frequency; very low quality evidence Acupuncture inferior to hormone therapy in QoL and frequency; no difference in severity; low quality evidence Chen 2016 (SR of 12 RCTs) (152): Seems to be effective; insufficient evidence; low quality evidence	‘Promising’; very low to moderate quality evidence <b>Upgrade to potential positive</b>
Nausea in pregnancy	<b>Matthews 2015 (Cochrane SR of interventions for nausea in pregnancy; 2 RCTs on acupuncture) (122):</b> No significant difference between real and sham acupuncture; low quality evidence	<b>No evidence of effect</b>
Hyperemesis gravidarum	Boelig 2016 (SR of interventions - 1 acupuncture study)(203): Insufficient evidence to identify clear differences between acupuncture and metoclopramide; very low quality evidence	<b>Unclear</b>
Melasma	Chai 2015 (SR of 8 RCTs; 2 high quality studies; 6 low to moderate)(204): Acupuncture appeared to be effective and safe; insufficient evidence	<b>Unclear</b>

**Table 6. All conditions reviewed sorted by clinical areas (contd)**

Condition	THE ACUPUNCTURE EVIDENCE PROJECT	Comments
<b>Gynaecology and obstetrics (contd)</b>		
Perimenopausal and postmenopausal sleep disturbance	Chiu 2016 (SR of 34 RCTs; 4 high quality)(138): Significant reduction in sleep disturbance which appears to be associated with changes in serum estradiol, FSH and LH; acupuncture recommended as adjunctive therapy in improving sleep disturbances in perimenopausal and postmenopausal women Bezerra 2015 (SR of 7 RCTs; 4 high quality; no studies with high risk of bias)(139): Improved sleep quality; limited evidence; moderate to high quality evidence	High quality studies favour acupuncture as adjunctive therapy; moderate to high quality evidence <b>Potential positive</b>
Assisted conception in ART	Manheimer 2013 (SR & MA of 16 RCTs) (205): Insufficient evidence <b>Cheong 2013 (Cochrane SR of 20 RCTs)(117):</b> No evidence that acupuncture improves live birth or pregnancy rates in ART regardless of whether performed around the time of oocyte retrieval or embryo transfer Shen 2015 (SR & MA)(118): Acupuncture ineffective when used only on the day of oocyte retrieval but effective when used at follicle phase and 25 min before and after embryo transfer Qian 2016 (SR of 30 RCTs & MA)(119) Acupuncture improves clinical pregnancy rates in women undergoing IVF. Optimal positive effects were seen when acupuncture was used during controlled ovarian hyperstimulation. Jo 2017 (SR of 4 RCTs & MA)(120) Acupuncture may increase the clinical pregnancy rate and ongoing pregnancy rate and decrease the risk of ovarian hyperstimulation syndrome in women with PCOS undergoing IVF or intracytoplasmic sperm injection	<b>Unclear</b>
Oocyte retrieval pain relief	<b>Kwan 2013 (Cochrane SR of 21 RCTs)(206):</b> Insufficient evidence; low quality evidence	<b>Unclear</b> (Cochrane update)
Polycystic ovarian syndrome	Ren 2014 (SR & MA of 31 studies)(207): Acupuncture may be effective; low quality evidence <b>Lim 2016 (Cochrane SR of 5 RCTs)(208):</b> Insufficient evidence; low to very low quality evidence	<b>Unclear</b>
Premenstrual syndrome	A SR review in 2014 included only studies before 2012, hence no new evidence (Jang 2014)(115) Insufficient evidence (Hofmeister 2016 - Narrative review)(116)	<b>Unclear</b> in USVA review 2014; no change in level
Primary ovarian insufficiency	Jo 2015 (SR of 8 RCTs & MA)(209): Acupuncture significantly lowered serum FSH levels and more women receiving acupuncture reported resumption of menses; low quality evidence	<b>Unclear</b>
Uterine fibroids	Dalton-Brewer 2016 (Narrative review of CAM)(105): Acupuncture may be a treatment option as part of a CAM treatment approach	<b>Unclear</b>

<b>Table 6. All conditions reviewed sorted by clinical areas (contd)</b>		
<b>Condition</b>	<b>THE ACUPUNCTURE EVIDENCE PROJECT</b>	<b>Comments</b>
<b>Gastrointestinal disorders</b>		
Constipation		<b>Potential positive</b> - USVA review 2014; no updates
Dyspepsia in diabetic gastroparesis (DGP)	Yang 2013 (SR of 14 RCTs)(210): May be effective for dyspepsia in DGP; insufficient evidence; low quality evidence	<b>Unclear</b>
Functional dyspepsia	<b>Lan 2014 (Cochrane SR of X RCTs)(211):</b> Insufficient evidence; low quality evidence Kim 2015 (SR of 20 RCTs; high risk of bias)(212): Acupuncture significantly superior to sham and medication; low quality evidence	<b>Unclear</b>
Inflammatory bowel disease	Langhorst 2015 (SR of 2 RCTs: one on Crohn's disease, one on ulcerative colitis)(213): Acupuncture superior to sham in disease activity and wellbeing but no different in QoL; low quality evidence (2 studies with low risk of bias but small samples)	<b>Unclear</b>
Irritable bowel syndrome	MacPherson 2016 (High quality RCT)(169): Acupuncture superior to usual care on IBS symptom severity score at 6, 9 and 12 months but not at 24 months	Unclear in USVA review 2014; <b>Upgrade to potential positive</b> - positive high quality RCT in 2016
Obesity	Esteghamati 2015 (Critical review of 3 SRs)(98): Acupuncture superior to medication, sham and lifestyle modification; low to very low quality evidence	<b>Potential positive</b> - USVA review 2014; no updates
<b>Headache and migraine</b>		
Headache	Kim 2012(76): Acupuncture is cost effective for dysmenorrhoea, allergic rhinitis, osteoarthritis & headache Coeytaux 2016 (Brief review of selected SRs and MAs)(44): A potentially important role for acupuncture as part of a treatment plan for migraine, tension-type headache, and several different types of chronic headache disorders. Cost-effective in Germany and UK Lardon 2016 (SR of 15 RCTs)(124): Acupuncture can reduce workplace headache pain intensity, frequency and related disability; low quality evidence	'A potentially important role for acupuncture' as part of a treatment plan for migraine, tension-type headache, and several different types of chronic headache disorders. <b>Positive</b>

**Table 6. All conditions reviewed sorted by clinical areas (contd)**

Condition	THE ACUPUNCTURE EVIDENCE PROJECT	Comments
<b>Headache and migraine (contd)</b>		
Migraines	<p>(Da Silva 2015 - Narrative review of large high quality RCTs)(40): Acupuncture seems to be at least as effective as conventional preventative medication for migraine and is safe, long lasting, and cost-effective</p> <p><b>Linde 2016 (Cochrane SR of 22 RCTs of at least 8 weeks duration)(41):</b> Acupuncture reduces migraine frequency. Acupuncture superior to sham and may be as effective as prophylactic drugs; subgroup analysis showed larger effect size when 16 or more treatments given (Z = 4.06) compared to 12 treatments or less (Z = 2.32); Moderate quality evidence</p> <p>Yang 2016 (SR of 10 RCTs)(42): Acupuncture superior to sham in effectiveness and reduced risk of recurrence; Moderate to high quality evidence</p>	Moderate to high quality evidence, safe and cost-effective (including Cochrane update); 16 or more treatments more effective than 12 treatments or less - <b>Positive</b>
Tension-type headache	<b>Linde 2016 (Cochrane SR of 12 RCTs)(43):</b> Effective for frequent episodic or chronic tension-type headaches; Moderate or low quality evidence	<b>Positive</b>
<b>Genitourinary/sexual disorders</b>		
Chronic kidney disease	<b>Kim 2016 (Cochrane SR of 24 RCTs or quasi-randomised CTs)(214):</b> Insufficient evidence; low to very low quality evidence	<b>Unclear</b>
Erectile dysfunction	Cui 2016 (SR of 3 RCTs): Insufficient evidence; low quality evidence	<b>Unclear</b>
Poor sperm quality	Jerng 2014 (SR of 4 RCTs)(215): Insufficient evidence; low quality evidence	<b>Unclear</b>
Prostatitis pain/chronic pelvic pain syndrome	<p>Chang 2016 (SR of 7 RCTs: 3 high quality, 1 moderate and 3 low)(96): Acupuncture superior to both sham and to usual care and safe, thus it should be offered when available</p> <p>Qin 2016 (SR of 7 RCTs)(97): Acupuncture superior to sham in pain, voiding and QoL; acupuncture superior to medication in pain relief but no different in voiding and QoL; acupuncture as an adjunctive treatment for symptom control should be considered; (3 high quality studies, 1 moderate and 3 low)</p>	<b>Potential positive</b>
Stress urinary incontinence in adults	<b>Wang 2013 (Cochrane SR of 1 study: acupuncture vs midodrine)(216):</b> Insufficient evidence; low quality evidence	<b>Unclear</b>
Urinary incontinence	Paik 2013 (SR of 4 RCTs)(217): Limited support for acupuncture or acupressure; Insufficient evidence; low to very low quality evidence	<b>Unclear</b>

**Table 6. All conditions reviewed sorted by clinical areas (contd)**

Condition	THE ACUPUNCTURE EVIDENCE PROJECT	Comments
<b>Surgery</b>		
Surgical conditions	Lee 2014 (Overview of 12 SRs on postsurgical nausea and vomiting and postsurgical pain)(163): Conflicting evidence; insufficient evidence	Generic systematic reviews on surgical conditions need to be more targeted to specific conditions
Ambulatory anaesthesia	Liodden 2013 (Narrative review)(165): Acupuncture may reduce preoperative anxiety, and postoperative pain, nausea, vomiting, shivering and emergence delirium. Acupuncture is safe and cost-effective. Acupuncture may be a beneficial adjunctive therapy for ambulatory anaesthesia.	Acupuncture safe, cost-effective and effective as an adjunctive therapy; no assessment of quality of evidence <b>Potential positive</b>
Postoperative nausea & vomiting	Cheong 2013 (SR of 30 RCTs)(56): Acupuncture and acupressure reduced number of cases of early nausea and vomiting (up to 24 hours); low quality evidence <b>Lee 2015 (Cochrane SR of 59 RCTs of PC 6 stimulation)(16):</b> PC 6 stimulation was superior to sham (low quality evidence); no difference between PC 6 stimulation and anti-emetics (moderate quality evidence); insufficient evidence that PC 6 plus anti-emetics is superior to anti-emetics alone.	<b>Upgrade to positive</b> (Cochrane update)
Postoperative nausea and vomiting and pain after tonsillectomy	Cho 2016 (MA of 12 RCTs; 7 high quality)(125): Pain scores, analgesic need and nausea and vomiting were reduced (acupuncture vs control); no significant adverse events; insufficient evidence Shin 2016 (SR & MA - 4 RCTs – 3 with low risk of bias; 3 randomised prospective studies and 1 pilot)(126): Acupuncture superior to controls and cost-effective; insufficient evidence	<b>Upgrade to positive</b> Cost effective
Postoperative pain	Wu 2016 (SR & MA of 13 RCTs)(17): Some forms of acupuncture (acupuncture, EA and TEAS) improved pain on day 1 after surgery and reduced opioid use; subgroup analysis showed acupuncture and TEAS superior to EA; moderate quality evidence	<b>Upgrade to positive</b>
Postoperative pain – back surgery	Cho 2015 (SR of 5 RCTs – 3 high quality)(57): Encouraging but limited evidence	<b>Upgrade to positive</b>
Postoperative pain – knee surgery	Barlow 2013 (SR of 5 RCTs: 3 acupuncture; 1 acupressure)(58): Acupressure reduced pain; acupuncture did not reduce pain but resulted in reduced use of ibuprofen; low quality evidence Chen 2015 (RCT acupuncture vs sham, n=60)(59): Acupuncture superior to sham in post-operative fentanyl use, time to first request for fentanyl and pain intensity	<b>Upgrade to positive</b>

**Table 6. All conditions reviewed sorted by clinical areas (contd)**

Condition	THE ACUPUNCTURE EVIDENCE PROJECT	Comments
<b>Surgery (contd)</b>		
Craniotomy anaesthesia	Asmussen 2016 (MA of 10 RCTs)(164): Acupuncture significantly reduced the amount of volatile anaesthetics and led to faster extubation time and postoperative patient recovery and reduced postoperative nausea and vomiting. In addition, significantly reduced blood levels of the brain tissue injury marker S100beta 48 hours after operation; (low quality studies excluded from MA)	MA which excluded low quality studies found positive effects; moderate to high quality evidence, but not definitive - <b>Potential positive</b>
Dentistry	Naik 2014 (Narrative review of 40 RCTs)(218): Acupuncture effective for inducing dental analgesia, relieving dental pain, myofascial pain and TMJ pain, controlling gag reflex, reducing dental anxiety and reducing post-operative pain and inflammation; no assessment of quality of evidence	'Promising' narrative review; no assessment of quality of evidence <b>Unclear</b>
Postoperative gastroparesis syndrome (PGS)	Cheong 2014 (SR of 16 RCTs; MA of 7)(219): Might be effective; insufficient evidence; low quality evidence	Insufficient evidence; low quality evidence <b>Unclear</b>
Postoperative ileus	Cheong 2016 (SR of 8 RCTs; MA of 4)(220): Acupuncture might be effective in reducing postoperative ileus; low quality evidence	<b>Unclear</b>
<b>Oncology</b>		
Aromatase-inhibitor-induced arthralgia	Bae 2015 (SR of 4 RCTs)(158): Acupuncture superior to sham in 2 high quality studies; no different from sham in 2 low-quality studies Chien 2015 (SR of 5 RCTs)(159): Acupuncture reduces joint pain and stiffness but not superior to sham; moderate to high quality evidence	<b>Potential positive</b>
Cancer-related insomnia	Choi 2016 (SR of 6 RCTs)(149): Acupuncture may be superior to sham acupuncture, drugs or hormones therapy. Number of studies and effect size are small for clinical significance; low quality evidence	<b>Unclear</b>
Cancer pain	<b>Pailey 2015 (Cochrane SR of 5 RCTs)(86):</b> Insufficient evidence; low to very low quality evidence Hu 2016 (SR of 20 RCTs)(221): Acupuncture plus medication superior to medication alone; very low quality evidence Chiu 2016 (SR of 29 RCTs)(87): Acupuncture relieved malignancy-related and surgery-induced pain but not pain induced by chemotherapy, radiotherapy or hormone therapy	Conflicting evidence <b>Unclear</b>

**Table 6. All conditions reviewed sorted by clinical areas (contd)**

Condition	THE ACUPUNCTURE EVIDENCE PROJECT	Comments
<b>Oncology (contd)</b>		
Cancer-related fatigue	<p>He 2013 (SR of 7 RCTs)(144): Acupuncture and moxibustion appear to be efficacious adjunctive therapy; Insufficient evidence; low quality evidence</p> <p>Ling 2013 (SR of RCTs)(145): Acupuncture and acupressure tend to be effective, acupuncture more than acupressure; low quality evidence</p> <p>Posadzki 2013 (SR of 7 RCTs)(146): Conflicting evidence: 4 studies showed acupuncture or acupuncture plus usual care superior to sham, usual care, enhanced usual care or no treatment; 3 studies showed no difference between acupuncture and sham; very low quality evidence</p> <p>Finnegan-John 2013 (SR of CAM 20 studies; 3 acupuncture/acupressure RCTs)(147): Acupuncture may reduce fatigue after cancer treatment; low quality evidence</p> <p>Zeng 2014 (MA of 7 studies)(148): Acupuncture plus education superior to usual care; low quality evidence</p>	<i>Potential positive</i>
Cancer-related psychological symptoms	Haddad 2014 (SR of 12 studies; 8 RCTs )(162): All included studies suggest benefits in depression, anxiety, sleep disturbance, and for improving QoL; strong evidence for safety; no assessment of quality of evidence	<i>Unclear</i>
Chemotherapy-induced nausea & vomiting (CINV)	<p>McKeon 2013 (SR of 7 acupuncture, 6 acupressure RCTs)(21): Acupuncture reduced the frequency of acute vomiting and the dose of rescue medication but did not reduce acute nausea severity or frequency compared to control. Acupressure showed a decrease in frequency of nausea but not acute vomiting or delayed symptoms. All studies used state-of-the-art combination antiemetics. Insufficient evidence due to underpowered studies; acupuncture low to moderate quality evidence; acupressure moderate to high quality evidence</p> <p>Garcia 2014 (SR update: 18 new RCTs)(55): Acupuncture is an appropriate referral option for chemotherapy-induced nausea and vomiting</p>	<i>Positive</i>
Chemotherapy-induced peripheral neuropathy	Franconi 2013 (SR of 3 RCTs, 3 case series, 1 rat study)(160): Acupuncture superior to sham in one RCT; very low quality evidence	<i>Unclear</i>
Myelosuppression after chemotherapy	Fu 2015 (Narrative review of 7 RCTs)(161): Insufficient evidence; low to very low quality evidence	<i>Unclear</i>
Recovery after colorectal cancer resection	Kim 2016 (SR of 7 RCTs)(157): Acupuncture efficacious and effective; Low to moderate quality evidence	<i>Potential positive</i>

**Table 6. All conditions reviewed sorted by clinical areas (contd)**

Condition	THE ACUPUNCTURE EVIDENCE PROJECT	Comments
<b>Oncology (contd)</b>		
Hot flushes in breast cancer	<p>Garcia 2015 (SR of 8 RCTs)(150): Acupuncture superior to sham in some studies and superior to baseline in all studies; low quality evidence</p> <p>Johns 2016: (SR of interventions; 2 acupuncture vs medication studies)(151): Acupuncture had similar efficacy to venlafaxine and gabapentin but may have longer durability after completing treatment and fewer side effects</p> <p>Chen 2016 (SR of 12 RCTs)(152): Acupuncture superior to sham in 3 studies; no different from sham in 6 studies; inferior to hormone therapy in 2 studies; low quality evidence</p> <p>Salehi 2016 (SR of 12 studies)(153): Conflicting evidence; low quality evidence</p> <p>Chiu 2016 (SR of 7 studies; 4 high quality)(87): Acupuncture yielded small-size effects on reducing hot-flash frequency and the severity of menopause-related symptoms</p>	<p>Insufficient or conflicting evidence; Low to high quality studies <b>Unclear</b></p>
Xerostomia in cancer	<p>Zhuang 2013 (SR of 4 studies)(154): Insufficient evidence</p> <p><b>Furness 2013 (Cochrane SR of non-pharmacological interventions 9 RCTs; 5 acupuncture)(155):</b> Small increase in saliva production; Low quality evidence</p> <p>Hanchanale 2015 (SR of 6 RCTs)(156): Acupuncture was superior to sham; low quality evidence</p>	<p><b>Unclear</b></p>
<b>Eye, ear, nose, throat</b>		
Allergic rhinitis	<p>Feng 2015 (SR of 13 RCTs)(54): Significant improvements in nasal symptoms scores and RQLQ; Moderate quality evidence</p> <p>Taw 2015 (SR of 2 large multi-centre RCTs, 3 acupuncture vs medication RCTs and 1 cost-effectiveness study)(15): High quality evidence of efficacy and effectiveness</p> <p>Seidman 2015 (OHNSF clinical practice guideline)(37): Option 5: Clinicians may offer acupuncture, or refer to a clinician who can offer acupuncture, for patients with AR who are interested in non-pharmacologic therapy; Aggregate evidence quality - Grade B</p> <p>McDonald 2016 (High quality RCT)(32): PAR - Significant improvements in symptoms, QoL, and reductions in total IgE and dust mite specific IgE and Substance P; high quality evidence</p> <p>Xue 2015 (High quality RCT)(222): SAR - Significant improvement in symptoms; high quality evidence</p> <p>Kim 2012(76): Acupuncture is cost effective for dysmenorrhoea, allergic rhinitis, osteoarthritis &amp; headache</p> <p>Reinhold 2103(77): SAR - Acupuncture significantly superior to rescue medication in QALY gained, but may cost more short term</p>	<p>Moderate to high quality evidence; safe and cost-effective for both SAR and PAR <b>Evidence of positive effect</b></p>
Dry eye	<p>Yang 2015 (SR of 7 RCTs)(166): Acupuncture was superior to artificial tears in tear break-up time, Schirmer I test, and cornea fluorescein staining; Low to moderate quality evidence</p>	<p><b>Upgrade from unclear to potential positive effect</b></p>

<b>Table 6. All conditions reviewed sorted by clinical areas (contd)</b>		
<b>Condition</b>	<b>THE ACUPUNCTURE EVIDENCE PROJECT</b>	<b>Comments</b>
<b>Eye, ear, nose, throat (contd)</b>		
Glaucoma	<b>Law 2013 (Cochrane SR; 1 RCT on auricular acupressure)(104):</b> Insufficient evidence; low quality evidence	<i>Unclear</i>
Meniere's disease/syndrome	He 2016 (SR of 12 RCTs)(223): Acupoint stimulation significantly effective in controlling vertigo but did not improve hearing loss; very low evidence	<i>Unclear</i>
Sudden sensorineural hearing loss	Zhang 2015 (SR of 12 RCTs)(224): Acupuncture plus usual care superior to usual care alone; low to very low quality evidence	<i>Unclear</i>
Tinnitus	He 2016 (SR of 5 RCTs)(91): EA - Insufficient evidence; low quality evidence	<i>Unclear</i>
<b>Paediatrics</b>		
Attention Deficit Hyperactivity Disorder (ADHD)	<b>Li 2011: (Cochrane SR; No eligible studies)(225):</b> Insufficient evidence	<i>Unclear</i> [not included in previous reviews]
Autism spectrum disorder (ASD)	<b>Cheuk 2011 (Cochrane SR of 10 RCTs)(226):</b> (4 high quality studies; 6 low quality) Insufficient evidence; low quality evidence	<i>Unclear</i> [not included in previous reviews]
Hypoxic ischemic encephalopathy in neonates	<b>Wong 2013 (Cochrane SR; No eligible studies)(227):</b> Insufficient evidence	<i>Unclear</i> [not included in previous reviews]
Mumps in children	<b>He 2015 (Cochrane SR; No eligible studies)(228):</b> Insufficient evidence	<i>Unclear</i> [not included in previous reviews]
Slowing progression of myopia	<b>Wei 2011 (Cochrane SR of 2 RCTs):</b> (2 high quality studies) Insufficient evidence; low quality evidence	<i>Unclear</i> [not included in previous reviews]
<b>Dermatology</b>		
Dermatology	Ma 2015 (SR of 15 RCTs: 3 Level I RCTs; 12 Level II)(229): Acupuncture improves outcome measures in the treatment of dermatitis, chloasma, pruritus, urticaria, hyperhidrosis, and facial elasticity	<i>Unclear</i>
Atopic dermatitis	Tan 2015 (SR: no eligible RCTs)(230): Insufficient evidence Vieira 2016 (Evidence-based review of CAM)(231): Acupuncture and acupressure 'promising therapies'; insufficient evidence	<i>Unclear</i>
Chronic urticaria	Yao 2015 (SR of 6 RCTs)(232): Acupuncture might be effective and safe for relieving symptoms of chronic urticaria; low level evidence	<i>Unclear</i>

<b>Table 6. All conditions reviewed sorted by clinical areas (contd)</b>		
<b>Condition</b>	<b>THE ACUPUNCTURE EVIDENCE PROJECT</b>	<b>Comments</b>
<b>Dermatology (contd)</b>		
Itch	Yu 2015 (SR & MA of 3 RCTs)(233): Acupuncture superior to sham and no treatment in alleviating itch; acupuncture might be effective for treating itch; insufficient evidence	<i>Unclear</i>
Psoriasis vulgaris	Coyle 2015 (SR of 6 RCTs)(234): Some evidence of benefit but also conflicting evidence; low quality evidence	<i>Unclear</i>
<b>Miscellaneous conditions</b>		
Acupuncture in Emergency Department	Kim 2013 (SR of 2 RCTs & 2 observational studies)(235): Insufficient evidence; low quality evidence	<i>Unclear</i>
Exercise performance & post-exercise recovery	USVA review 2014 'unclear'; no updates	<i>Unclear</i>
Fatigue in systemic lupus erythematosus	del Pino-Sedeno 2016 (SR of non-pharmacological interventions; 1 acupuncture study)(236): Acupuncture and minimal acupuncture superior to usual care; moderate quality study	<i>Unclear</i>
Primary Sjogren's syndrome	Hackett 2015 (SR of non-pharmacological interventions; 1 acupuncture study)(237): Acupuncture not superior to usual care in increasing salivary flow rate; moderate quality study	<i>Unclear</i>
Sensory perception	Baeumler 2014 (MA of 85 high quality studies)(238): Acupuncture effects sensory thresholds especially pressure pain threshold; high quality evidence	<i>Potential positive effect</i>

**Table 7. Conditions reported in this review with evidence of cost-effectiveness**

Condition	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)	Comments
Allergic Rhinitis	Taw 2015 (SR of 2 large multi-centre RCTs, 3 acupuncture vs medication RCTs and 1 cost-effectiveness study) Kim 2012(Cost effectiveness analysis)(76): Acupuncture is cost effective for dysmenorrhoea, allergic rhinitis, osteoarthritis & headache.	Safe and <b>cost-effective</b> for both SAR and PAR.
Ambulatory Anaesthesia	Liodden 2013 (Narrative review)(165): Acupuncture may reduce preoperative anxiety, and postoperative pain, nausea, vomiting, shivering and emergence delirium. Acupuncture is safe and cost-effective. Acupuncture may be a beneficial adjunctive therapy for ambulatory anaesthesia.	Acupuncture safe, <b>cost-effective</b> and effective as an adjunctive therapy.
Chronic Pain	MacPherson Oct 2016 (SR & MA of 29 trials) (239)The effects of a course of acupuncture treatment for patients with chronic pain do not appear to decrease importantly over 12 months. Patients can generally be reassured that treatment effects persist. Studies of the cost-effectiveness of acupuncture should take our findings into account.	<b>“Studies of the cost-effectiveness of acupuncture should take our findings into account.”</b>
Depression	Spackman 2014 (Cost-effectiveness analysis)(89): Acupuncture is cost-effective compared with counselling or usual care alone, although the ranking of counselling and acupuncture depends on the relative cost of delivering these interventions.	<b>Cost-effective</b>
Dysmenorrhoea	Kim 2012 (Cost effectiveness analysis)(76): Acupuncture is cost effective for dysmenorrhoea, allergic rhinitis, osteoarthritis & headache.	<b>Cost effective</b>
Headache	Kim 2012 (Cost effectiveness analysis)(76): Acupuncture is cost effective for dysmenorrhoea, allergic rhinitis, osteoarthritis & headache. Coeytaux 2016 (Brief review of selected SRs and MAs)(44): A potentially important role for acupuncture as part of a treatment plan for migraine, tension-type headache, and several different types of chronic headache disorders. Cost-effective in Germany and UK.	A potentially important role for acupuncture’ as part of a treatment plan for migraine, tension-type headache, and several different types of chronic headache disorders. <b>Cost effective</b>
Low back pain	Taylor 2014 (Cost effectiveness analysis/MA)(51): Cost effective for chronic low back pain. Andronis 2016 (SR of 33 studies)(50): Likely to be cost effective.	Moderate to high quality evidence <b>Cost effective, safe.</b>
Migraine	Da Silva 2015 (Narrative review of large high quality RCTs)(40): Acupuncture seems to be at least as effective as conventional preventative medication for migraine and is safe, long lasting, and cost-effective.	Moderate to high quality evidence, safe and <b>cost-effective</b> (including Cochrane update); 16 or more treatments more effective than 12 treatments or less.
Neck Pain	Van der Velde 2015 (SR of 6 studies)(175): Acupuncture plus usual medical care is cost-effective for neck pain and its associated disorders (NAD).	Moderate quality evidence (Cochrane update); Acupuncture plus medication is <b>cost-effective.</b>
Osteoarthritis	Kim 2012 (Cost effectiveness analysis)(76): Acupuncture is cost effective for dysmenorrhoea, allergic rhinitis, osteoarthritis & headache.	<b>Cost-effective</b>
Post-operative nausea and vomiting	Shin 2016 (SR & MA - 4 RCTs – 3 with low risk of bias; 3 randomised prospective studies and 1 pilot)(126): Acupuncture superior to controls and cost-effective.	<b>Cost effective</b>

**Table 8. Conditions reported in this review with evidence of safety**

Condition	The Acupuncture Evidence Project (Mar 2013 - Sept 2016)	Comments
Acupuncture generally prior to this review	Zhang et al 2010 (Review of 98 case reports and 17 case series)(240) ‘Various types of acupuncture-related adverse events have been reported in China. Similar events have been reported by other countries, usually as a result of inappropriate technique. Acupuncture can be considered inherently safe in the hands of well-trained practitioners.’	<b>Acupuncture can be considered inherently safe in the hands of well-trained practitioners.</b>
Allergic Rhinitis	Taw 2015 (SR of 2 large multi-centre RCTs, 3 acupuncture vs medication RCTs and 1 cost-effectiveness study). Kim 2012(76): Acupuncture is cost effective for dysmenorrhoea, allergic rhinitis, osteoarthritis & headache.	<b>Safe</b> and cost-effective for both SAR and PAR.
Ambulatory Anaesthesia	Liudden 2013 (Narrative review)(165): Acupuncture may reduce preoperative anxiety, and postoperative pain, nausea, vomiting, shivering and emergence delirium. Acupuncture is safe and cost-effective. Acupuncture may be a beneficial adjunctive therapy for ambulatory anaesthesia.	Acupuncture <b>safe</b> , cost-effective and effective as an adjunctive therapy.
Alzheimer's disease	Zhou 2015 (SR & MA of 10 RCTs)(200): Acupuncture superior to medication in improving cognitive function on MMSE. Acupuncture plus medication superior to medication alone. Acupuncture is safe.	Acupuncture is <b>Safe</b> .
Cancer-related psychological symptoms	Haddad 2014 (SR of 12 studies; 8 RCTs)(162): All included studies suggest benefits in depression, anxiety, sleep disturbance, and for improving QoL; strong evidence for safety; no assessment of quality of evidence.	Strong evidence for <b>safety</b> .
Depression	Chan 2015 (SR & MA of 13 RCTs;1 high quality, 5 moderate, 7 low)(88): Acupuncture plus SSRIs superior to SSRIs alone, with an early onset of action and was safe and well-tolerated; EA had greater effect than manual acupuncture. Bosch 2015 (Review of SRs & MAs)(82): Effective and safe for major depressive disorder, especially in improving sleep, mood and QoL by modulating and normalizing the limbic–paralimbic–neocortical network (LPNN), including the default mode network (DMN); ‘promising’ evidence.	<b>Safe</b> and well tolerated. Effective and <b>safe</b> for major depressive disorder.
Low back pain	Nahin 2016 (4 RCTs; Excluded studies not performed in USA or by US researchers)(195): Acupuncture superior to usual care; Acupuncture superior to sham in 1 RCT, but not superior in 2 RCTs. NIH (2016) Promise in the following for safety and effectiveness in treating pain: Acupuncture and yoga for back pain, acupuncture and tai chi for osteoarthritis of the knee (241). Chou et al 2016 (Comparative effectiveness review) (47): Serious adverse events were not reported in any trial.	Moderate to high quality evidence Cost effective. Promise in <b>safety</b> and effectiveness. Serious adverse events were not reported in any trial.
Migraine	Da Silva 2015 (Narrative review of large high quality RCTs)(40): Acupuncture seems to be at least as effective as conventional preventative medication for migraine and is safe, long lasting, and cost-effective.	Moderate to high quality evidence, <b>safe</b> and cost-effective (including Cochrane update); 16 or more treatments more effective than 12 treatments or less.
Osteoarthritis of the Knee	Nahin 2016 (4 RCTs; Excluded studies not performed in USA or by US researchers)(195). NIH 2016 (241): Promise in the following for safety and effectiveness in treating pain: Acupuncture and yoga for back pain, acupuncture and tai chi for osteoarthritis of the knee.	Promise in <b>safety</b> and effectiveness.
Prostatitis pain/chronic pelvic pain syndrome	Chang 2016 (SR of 7 RCTs: 3 high quality studies, 1 moderate and 3 low)(96)Acupuncture superior to both sham and to usual care and safe, thus it should be offered when available.	Acupuncture superior to both sham and to usual care and <b>safe</b> .

## APPENDIX

**Table 9. NHMRC Evidence Hierarchy: designations of 'levels of evidence' according to type of research question (242)**

Level	Intervention	Diagnostic accuracy	Prognosis	Aetiology	Screening Intervention
I	A systematic review of level II studies	A systematic review of level II studies	A systematic review of level II studies	A systematic review of level II studies	A systematic review of level II studies
II	A randomised controlled trial	A study of test accuracy with: an independent, blinded comparison with a valid reference standard, among consecutive persons with a defined clinical presentation	A prospective cohort study	A prospective cohort study	A randomised controlled trial
III-1	A pseudorandomised controlled trial (i.e. alternate allocation or some other method)	A study of test accuracy with: an independent, blinded comparison with a valid reference standard, among non-consecutive persons with a defined clinical presentation	All or none	All or none	A pseudorandomised controlled trial (i.e. alternate allocation or some other method)
III-2	A comparative study with concurrent controls: <ul style="list-style-type: none"> <li>▪ Non-randomised, experimental trial</li> <li>▪ Cohort study</li> <li>▪ Case-control study</li> <li>▪ Interrupted time series with a control group</li> </ul>	A comparison with reference standard that does not meet the criteria required for Level II and III-1 evidence	Analysis of prognostic factors amongst persons in a single arm of a randomised controlled trial	A retrospective cohort study	A comparative study with concurrent controls: <ul style="list-style-type: none"> <li>▪ Non-randomised, experimental trial</li> <li>▪ Cohort study</li> <li>▪ Case-control study</li> </ul>
III-3	A comparative study without concurrent controls: <ul style="list-style-type: none"> <li>▪ Historical control study</li> <li>▪ Two or more single arm study</li> <li>▪ Interrupted time series without a parallel control group</li> </ul>	Diagnostic case-control study	A retrospective cohort study	A case-control study	A comparative study without concurrent controls: <ul style="list-style-type: none"> <li>▪ Historical control study</li> <li>▪ Two or more single arm study</li> </ul>
IV	Case series with either post-test or pre-test/post-test outcomes	Study of diagnostic yield (no reference standard)	Case series, or cohort study of persons at different stages of disease	A cross-sectional study or case series	Case series

## GRADE and guideline development

GRADE quality levels reflect how much confidence the reviewers have that the estimate of effect is close to the true effect in a systematic review. GRADE considers five factors in making this judgement: imprecision, inconsistency, indirectness of study results, publication bias and bias generally. GRADE levels do not necessarily infer a recommendation, and the GRADE process is separate to the process of making recommendations. Although a high level of evidence is likely to lead to a recommendation, low or very low evidence can lead to a strong recommendation in some cases. The development of recommendations involves more than just the quality of evidence and requires consideration of other factors including cost, clinical judgement and patient preference (8).

Quality level	Definition
High	We are very confident that the true effect lies close to that of the estimate of the effect
Moderate	We are moderately confident in the effect estimate: The true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different
Low	Our confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect
Very low	We have very little confidence in the effect estimate: The true effect is likely to be substantially different from the estimate of effect

## PLAIN ENGLISH SUMMARY

### Bottom Line

Our study found evidence for the effectiveness of acupuncture for 117 conditions, with stronger evidence for acupuncture's effectiveness for some conditions than others. Acupuncture is considered safe in the hands of a well-trained practitioner and has been found to be cost effective for some conditions. The quality and quantity of research into acupuncture's effectiveness is increasing.

### Background

Acupuncture originated in China and is now practised throughout the world. Although acupuncture has been practised for thousands of years, evidence of its effectiveness is still controversial. The Australian Acupuncture and Chinese Medicine Association Ltd (AACMA) identified the need for an updated review of the evidence with greater rigour than was possible in the past and commissioned The Acupuncture Evidence Project.

We searched the literature with a focus on systematic reviews and meta analyses (the highest form of evidence available). We sorted the evidence to identify the conditions for which acupuncture has been found to be most effective for. We also looked for evidence of acupuncture's safety and cost-effectiveness, and we reported how the evidence for acupuncture's effectiveness has changed over an eleven-year time-frame.

### Key results

Of the 122 conditions identified, strong evidence supported the effectiveness of acupuncture for eight conditions, moderate evidence supported the use of acupuncture for a further 38 conditions, weak positive/unclear evidence supported the use of acupuncture for 71 conditions, and little or no evidence was found for the effectiveness of acupuncture for five conditions (meaning that further research is needed to clarify the effectiveness of acupuncture in these last two categories).

*It is no longer possible to say that the effectiveness of acupuncture is because of the placebo effect, or that it is useful only for musculoskeletal pain.*

In addition, research showed that acupuncture was cost effective for 10 conditions, and is safe in the hands of a well-trained practitioner. The level of evidence has increased over the 11-year period of this study for 24 conditions. Placebo-controlled clinical trials consistently underestimate the true effect size of acupuncture (which means that acupuncture is more effective than the type of trials used in this review show), yet they have still demonstrated National Health and Medical Research Council (NHMRC) Level I evidence for the effectiveness of acupuncture for 117 conditions.

## Summary of Findings

**Summary of Findings 1:** The following tables summarise the effectiveness of acupuncture for various conditions.

<b>Table 1. Conditions with strong evidence supporting the effectiveness of acupuncture</b>	
Reviews with consistent statistically significant positive effects and where authors have recommended the intervention. The quality of evidence is rated as moderate or high quality.	
<ul style="list-style-type: none"> <li>- Allergic rhinitis (perennial &amp; seasonal)</li> <li>- Chemotherapy-induced nausea and vomiting (with anti-emetics)</li> <li>- Chronic low back pain</li> <li>- Headache (tension-type and chronic)</li> </ul>	<ul style="list-style-type: none"> <li>- Knee osteoarthritis</li> <li>- Migraine prophylaxis</li> <li>- Postoperative nausea &amp; vomiting</li> <li>- Postoperative pain</li> </ul>

<b>Table 2. Conditions with moderate evidence supporting the effectiveness of acupuncture</b>	
Reviews reporting all individual RCTs or pooled effects across RCTs as positive, but the reviewers deeming the evidence insufficient to draw firm conclusions. The quality of evidence is rated as moderate or high quality.	
<ul style="list-style-type: none"> <li>- Acute low back pain</li> <li>- Acute stroke</li> <li>- Ambulatory anaesthesia</li> <li>- Anxiety</li> <li>- Aromatase-inhibitor-induced arthralgia</li> <li>- Asthma in adults</li> <li>- Back or pelvic pain during pregnancy</li> <li>- Cancer pain</li> <li>- Cancer-related fatigue</li> <li>- Constipation</li> <li>- Craniotomy anaesthesia</li> <li>- Depression (with antidepressants)</li> <li>- Dry eye</li> <li>- Hypertension (with medication)</li> <li>- Insomnia</li> <li>- Irritable bowel syndrome</li> <li>- Labour pain</li> <li>- Lateral elbow pain</li> <li>- Menopausal hot flushes</li> </ul>	<ul style="list-style-type: none"> <li>- Modulating sensory perception thresholds</li> <li>- Neck pain</li> <li>- Obesity</li> <li>- Perimenopausal &amp; postmenopausal insomnia</li> <li>- Plantar heel pain</li> <li>- Post-stroke insomnia</li> <li>- Post-stroke shoulder pain</li> <li>- Post-stroke spasticity</li> <li>- Post-traumatic stress disorder</li> <li>- Prostatitis pain/chronic pelvic pain syndrome</li> <li>- Recovery after colorectal cancer resection</li> <li>- Restless leg syndrome</li> <li>- Schizophrenia (with antipsychotics)</li> <li>- Sciatica</li> <li>- Shoulder impingement syndrome (early stage) (with exercise)</li> <li>- Shoulder pain</li> <li>- Smoking cessation (up to 3 months)</li> <li>- Stroke rehabilitation</li> <li>- Temporomandibular pain</li> </ul>

**Summary of Findings 1 (continued):** The following tables summarise the effectiveness of acupuncture for various conditions

<b>Table 3. Conditions with weak positive/unclear evidence supporting the effectiveness of acupuncture</b>	
Reviews consisted mostly of weak positive evidence or conflicting evidence between reviews or between authors within a review, with reviewers summarising the evidence as inconclusive. Reviews are of low or very low quality; or there is conflicting levels of evidence within or between reviews.	
<ul style="list-style-type: none"> <li>- Acupuncture in Emergency Department</li> <li>- Acute ankle sprain in adults</li> <li>- Alzheimer’s disease</li> <li>- Angina pectoris</li> <li>- Assisted conception in ART</li> <li>- Asthma in children</li> <li>- Atopic dermatitis</li> <li>- Attention Deficit Hyperactivity Disorder (ADHD)</li> <li>- Autism spectrum disorder (ASD)</li> <li>- Bell’s palsy</li> <li>- Bladder pain syndrome</li> <li>- Cancer-related insomnia</li> <li>- Cancer-related psychological symptoms</li> <li>- Carpal tunnel syndrome</li> <li>- Chemotherapy-induced peripheral neuropathy</li> <li>- Chronic fatigue syndrome</li> <li>- Chronic kidney disease</li> <li>- Chronic obstructive pulmonary disease (COPD)</li> <li>- Chronic urinary retention due to spinal cord injury</li> <li>- Chronic urticaria</li> <li>- Dysmenorrhoea</li> <li>- Dyspepsia in diabetic gastroparesis (DGP)</li> <li>- Erectile dysfunction</li> <li>- Exercise performance &amp; post-exercise recovery</li> <li>- Fatigue in systemic lupus erythematosus</li> <li>- Fibromyalgia</li> <li>- Functional dyspepsia</li> <li>- Gag reflex in dentistry</li> <li>- Glaucoma</li> <li>- Heart failure</li> <li>- Hot flushes in breast cancer</li> <li>- Hyperemesis gravidarum</li> <li>- Hypoxic ischemic encephalopathy in neonates</li> <li>- Induction of labour</li> <li>- Inflammatory bowel disease</li> </ul>	<ul style="list-style-type: none"> <li>- Itch</li> <li>- Lumbar spinal stenosis</li> <li>- Melasma</li> <li>- Meniere’s disease/syndrome</li> <li>- Menopausal syndrome</li> <li>- Multiple sclerosis</li> <li>- Mumps in children</li> <li>- Myelosuppression after chemotherapy</li> <li>- Oocyte retrieval pain relief</li> <li>- Opiate addiction</li> <li>- Opioid detoxification</li> <li>- Parkinson’s disease</li> <li>- Polycystic ovarian syndrome</li> <li>- Poor sperm quality</li> <li>- Postnatal depression</li> <li>- Postoperative gastroparesis syndrome (PGS)</li> <li>- Postoperative ileus</li> <li>- Post-stroke hiccoughs</li> <li>- Premenstrual syndrome</li> <li>- Primary ovarian insufficiency</li> <li>- Primary Sjogren’s syndrome</li> <li>- Psoriasis vulgaris</li> <li>- Rheumatoid arthritis Slowing progression of myopia</li> <li>- Spinal cord injury</li> <li>- Stress urinary incontinence in adults</li> <li>- Sudden sensorineural hearing loss</li> <li>- Surgery analgesia</li> <li>- Tinnitus</li> <li>- Traumatic brain injury</li> <li>- Urinary incontinence</li> <li>- Uterine fibroids</li> <li>- Vascular cognitive impairment without dementia</li> <li>- Vascular dementia</li> <li>- Whiplash associated disorder (WAD)</li> <li>- Xerostomia in cancer</li> </ul>

**Summary of Findings 1 (continued):** The following tables summarise the effectiveness of acupuncture for various conditions

<b>Table 4. Conditions with little or no evidence supporting the effectiveness of acupuncture</b>	
Reviews have consistently found little support for acupuncture. The quality of the evidence is consistently low or very low. Further research required.	
- Alcohol dependence	- Nausea in pregnancy
- Cocaine addiction	- Smoking cessation (more than 6 months)
- Epilepsy	

**Summary of Findings 2:** Conditions with evidence of cost-effectiveness.

<b>Table 5. Conditions with evidence of cost effectiveness</b>	
- Allergic Rhinitis	- Low back pain
- Ambulatory Anaesthesia	- Migraine
- Chronic Pain	- Neck Pain (plus usual medical care)
- Depression	- Osteoarthritis
- Dysmenorrhoea	- Post-operative nausea and vomiting
- Headache	

**Summary of Findings 3:** Conditions with evidence of safety.

<b>Table 6. Conditions with evidence of safety</b>	
Condition	Comments
Acupuncture generally prior to this review	Acupuncture can be considered inherently safe in the hands of well-trained practitioners.
Allergic Rhinitis	<b>Safe</b> and cost-effective
Ambulatory Anaesthesia	Acupuncture <b>safe</b> , cost-effective and effective as an adjunctive therapy.
Alzheimers disease	Acupuncture is <b>Safe</b> .
Cancer-related psychological symptoms	Strong evidence for <b>safety</b> .
Depression	Strong evidence for <b>safety</b> . Effective and <b>safe</b> for major depressive disorder.
Low back pain	<b>Safe</b> and well tolerated.
Migraine	Moderate to high quality evidence Cost effective. Promise in <b>safety</b> and effectiveness. Serious adverse events were not reported in any trial.
Osteoarthritis of the Knee	Promise in <b>safety</b> and effectiveness.
Prostatitis pain/chronic pelvic pain syndrome	Acupuncture superior to both sham and to usual care and <b>safe</b> .

**Summary of Findings 4:** Changes in evidence levels over the eleven-year period covered by this review

<b>Table 7. Statistical summary of findings of this review</b>			
<b>Evidence Level</b>	<b>Number of Conditions</b>	<b>Changes in Level of Evidence</b>	<b>Number of Conditions</b>
Strong Evidence of effect	8	Increase to strong evidence	5
Moderate Evidence effect	38	Increase to moderate evidence	18
Unclear/mixed evidence	71	Increase to weak positive/unclear evidence	1
Little of no evidence of effect	5	Decreased evidence level	2
Total conditions with some evidence of effect (any level)	117	_____	_____
Total conditions reviewed	122	Total increases in evidence level since prior reviews	24

## REFERENCES

1. Bannerman R. Acupuncture. The World Health Organization View. Geneva. The World Health Organisation; 1979.
2. Chmielnicki B. Evidence-Based Acupuncture: WHO official position. Chmielnicki, B.; 2014 [3 December, 2016]; Available from: <http://www.evidencebasedacupuncture.org/who-official-position/>.
3. NIH Consensus Conference. Acupuncture. JAMA. 1998 Nov 4;280(17):1518-24.
4. World Health Organization. Acupuncture: review and analysis of reports on controlled clinical trials. Geneva: World Health Organization; 2002.
5. Biotext. Alternative therapies and Department of Veterans' Affairs Gold and White Card arrangements. In: Australian Government Department of Veterans' Affairs, editor: Australian Government Department of Veterans' Affairs; 2010.
6. Hempel S, Taylor SL, Solloway MR, Miake-Lye IM, Beroes JM, Shanman R, et al. VA Evidence-based Synthesis Program Reports. Evidence Map of Acupuncture. Washington (DC): Department of Veterans Affairs; 2014.
7. National Health and Medical Research Council. NHMRC additional levels of evidence and grades for recommendations for developers of guidelines. National Health and Medical Research Council; 2009.
8. Balshem H, Helfand M, Schunemann HJ, Oxman AD, Kunz R, Brozek J, et al. GRADE guidelines: 3. Rating the quality of evidence. J Clin Epidemiol. 2011 Apr;64(4):401-6.
9. Wehling M. Non-steroidal anti-inflammatory drug use in chronic pain conditions with special emphasis on the elderly and patients with relevant comorbidities: management and mitigation of risks and adverse effects. Eur J Clin Pharmacol. 2014 Oct;70(10):1159-72.
10. Krashin D, Murinova N, Sullivan M. Challenges to Treatment of Chronic Pain and Addiction During the 'Opioid Crisis'. Curr Pain Headache Rep. 2016 Dec;20(12):65.
11. Sostres C, Gargallo CJ, Lanás A. Nonsteroidal anti-inflammatory drugs and upper and lower gastrointestinal mucosal damage. Arthritis Res Ther. 2013;15 Suppl 3:S3.
12. Schmidt M, Lamberts M, Olsen AM, Fosboll E, Niessner A, Tamargo J, et al. Cardiovascular safety of non-aspirin non-steroidal anti-inflammatory drugs: review and position paper by the working group for Cardiovascular Pharmacotherapy of the European Society of Cardiology. Eur Heart J Cardiovasc Pharmacother. 2016 Apr;2(2):108-18.
13. Fransen M, Nairn L, Bridgett L, Crosbie J, March L, Parker Mbbs D, et al. Post-acute rehabilitation after total knee replacement: A multicentre randomized clinical trial comparing long-term outcomes. Arthritis Care Res (Hoboken). 2016 Nov 21.
14. Steiner TJ, Stovner LJ, Birbeck GL. Migraine: the seventh disabling. J Headache Pain. 2013 Jan 10;14:1.
15. Taw MB, Reddy WD, Omole FS, Seidman MD. Acupuncture and allergic rhinitis. Curr Opin Otolaryngol Head Neck Surg. 2015 Jun;23(3):216-20.
16. Lee A, Chan SK, Fan LT. Stimulation of the wrist acupuncture point PC6 for preventing postoperative nausea and vomiting. Cochrane Database Syst Rev. 2015(11):Cd003281.
17. Wu MS, Chen KH, Chen IF, Huang SK, Tzeng PC, Yeh ML, et al. The Efficacy of Acupuncture in Post-Operative Pain Management: A Systematic Review and Meta-Analysis. PLOS ONE. 2016;11(3):e0150367.

18. Center for Disease Control and Prevention. Osteoarthritis. In: US Department of Health and Human Services, editor. USA. 2015.
19. Cross M, Smith E, Hoy D, Nolte S, Ackerman I, Fransen M, et al. The global burden of hip and knee osteoarthritis: estimates from the global burden of disease 2010 study. *Ann Rheum Dis.* 2014;73(7):1323-30.
20. Hoy D, March L, Brooks P, Blyth F, Woolf A, Bain C, et al. The global burden of low back pain: estimates from the Global Burden of Disease 2010 study. *Ann Rheum Dis.* 2014 Jun;73(6):968-74.
21. McKeon C, Smith, C.A., Hardy, J. & Chang, E. Acupunctrue and acupressure for chemotherapy-induced nausea and vomiting : a systematic review. *Australian Journal of Acupuncture and Chinese Medicine.* 2013;8(1):2-17.
22. Morris ZS, Wooding S, Grant J. The answer is 17 years, what is the question: understanding time lags in translational research. *J R Soc Med.* 2011 Dec;104(12):510-20.
23. Birch S. A review and analysis of placebo treatments, placebo effects, and placebo controls in trials of medical procedures when sham is not inert. *J Altern Complement Med.* 2006 Apr;12(3):303-10.
24. Linde K, Niemann K, Meissner K. Are sham acupuncture interventions more effective than (other) placebos? A re-analysis of data from the Cochrane review on placebo effects. *Forsch Komplementmed.* 2010 Oct;17(5):259-64.
25. Langevin HM, Wayne PM, Macpherson H, Schnyer R, Milley RM, Napadow V, et al. Paradoxes in acupuncture research: strategies for moving forward. *Evid Based Complement Alternat Med.* 2011;2011:180805.
26. Lund I, Lundeberg T. Are minimal, superficial or sham acupuncture procedures acceptable as inert placebo controls? *Acupunct Med.* 2006 Mar;24(1):13-5.
27. Lund I, Naslund J, Lundeberg T. Minimal acupuncture is not a valid placebo control in randomised controlled trials of acupuncture: a physiologist's perspective. *Chin Med.* 2009;4:1.
28. Linde K, Niemann K, Schneider A, Meissner K. How large are the nonspecific effects of acupuncture? A meta-analysis of randomized controlled trials. *BMC Med.* 2010;8:75.
29. Lundeberg T, Lund I, Sing A, Naslund J. Is placebo acupuncture what it is intended to be? *Evid Based Complement Alternat Med.* 2011;2011:932407.
30. Patsopoulos NA. A pragmatic view on pragmatic trials. *Dialogues Clin Neurosci.* 2011 Jun;13(2):217-24.
31. National Centre for Complementary and Alternative Health. Acupuncture Research - Areas of High and Low Programmatic Priorities. US Department of Health and Human Services; 2015 [cited 2016]; Available from: <https://nccih.nih.gov/grants/acupuncture/priorities?nav=fb>.
32. McDonald JL, Smith PK, Smith CA, Changli Xue C, Golianu B, Cripps AW. Effect of acupuncture on house dust mite specific IgE, substance P, and symptoms in persistent allergic rhinitis. *Ann Allergy Asthma Immunol.* 2016 Jun;116(6):497-505.
33. Schug SA, Palmer GM, Scott DA, Halliwell R, Trinca J. Acute pain management: scientific evidence, fourth edition, 2015. *Med J Aust.* 2016 May 2;204(8):315-7.
34. Hopman K KL, Lukersmith S, McColl AR, & Vine K. Clinical Practice Guidelines for the Management of Rotator Cuff Syndrome in the Workplace. The University of New South Wales. 2013.
35. Gan TJ, Diemunsch P, Habib AS, Kovac A, Kranke P, Meyer TA, et al. Consensus guidelines for the management of postoperative nausea and vomiting. *Anesth Analg.* 2014 Jan;118(1):85-113.

36. Birch S, Alraek T, Lee MS. Challenges for clinical practice guidelines in traditional medicines: The example of acupuncture. *European Journal of Integrative Medicine*. <http://dx.doi.org/doi:10.1016/j.eujim.2016.07.032>
37. Seidman MD, Gurgel RK, Lin SY, Schwartz SR, Baroody FM, Bonner JR, et al. Clinical practice guideline: Allergic rhinitis. *Otolaryngol Head Neck Surg*. 2015 Feb;152(1 Suppl):S1-43.
38. National Institute for Health Care and Excellence. Management of migraine (with or without aura): NICE guideline CG150. In: National Institute for Health Care and Excellence, editor. 2012 (updated 2015).
39. Scottish Incollegiate Guidelines Network (SIGN). Management of Chronic Pain (SIGN publication no. 136). Scottish Incollegiate Guidelines Network (SIGN); 2013.
40. Da Silva AN. Acupuncture for migraine prevention. *Headache*. 2015 Mar;55(3):470-3.
41. Linde K, Allais G, Brinkhaus B, Fei Y, Mehring M, Vertosick EA, et al. Acupuncture for the prevention of episodic migraine. *Cochrane Database Syst Rev*. 2016(6):Cd001218.
42. Yang Y, Que Q, Ye X, Zheng G. Verum versus sham manual acupuncture for migraine: a systematic review of randomised controlled trials. *Acupunct Med*. 2016 Apr;34(2):76-83.
43. Linde K, Allais G, Brinkhaus B, Fei Y, Mehring M, Shin BC, et al. Acupuncture for the prevention of tension-type headache. *Cochrane Database Syst Rev*. 2016;4:Cd007587.
44. Coeytaux RR, Befus D. Role of Acupuncture in the Treatment or Prevention of Migraine, Tension-Type Headache, or Chronic Headache Disorders. *Headache*. 2016 Jul;56(7):1238-40.
45. Chou R, Deyo R, Friedly J, Skelly A, Hashimoto R, Weimer M, et al. AHRQ Comparative Effectiveness Reviews. Noninvasive Treatments for Low Back Pain. Rockville (MD): Agency for Healthcare Research and Quality (US); 2016.
46. Lam M, Galvin R, Curry P. Effectiveness of acupuncture for nonspecific chronic low back pain: a systematic review and meta-analysis. *Spine (Phila Pa 1976)*. 2013 Nov 15;38(24):2124-38.
47. Lee JH, Choi TY, Lee MS, Lee H, Shin BC, Lee H. Acupuncture for acute low back pain: a systematic review. *Clin J Pain*. 2013 Feb;29(2):172-85.
48. Wellington J. Noninvasive and alternative management of chronic low back pain (efficacy and outcomes). *Neuromodulation*. 2014 Oct;17 Suppl 2:24-30.
49. Liu L, Skinner M, McDonough S, Mabire L, Baxter GD. Acupuncture for low back pain: an overview of systematic reviews. *Evid Based Complement Alternat Med*. 2015;2015:328196.
50. Andronis L, Kinghorn P, Qiao S, Whitehurst DG, Durrell S, McLeod H. Cost-Effectiveness of Non-Invasive and Non-Pharmacological Interventions for Low Back Pain: a Systematic Literature Review. *Appl Health Econ Health Policy*. 2016 Aug 22.
51. Taylor P, Pezzullo L, Grant SJ, Bensoussan A. Cost-effectiveness of Acupuncture for Chronic Nonspecific Low Back Pain. *Pain Pract*. 2014 Sep;14(7):599-606.
52. Corbett MS, Rice SJ, Madurasinghe V, Slack R, Fayter DA, Harden M, et al. Acupuncture and other physical treatments for the relief of pain due to osteoarthritis of the knee: network meta-analysis. *Osteoarthritis Cartilage*. 2013 Sep;21(9):1290-8.
53. Manyanga T, Froese M, Zarychanski R, Abou-Setta A, Friesen C, Tennenhouse M, et al. Pain management with acupuncture in osteoarthritis: a systematic review and meta-analysis. *BMC Complement Altern Med*. 2014;14:312.

54. Feng S, Han M, Fan Y, Yang G, Liao Z, Liao W, et al. Acupuncture for the treatment of allergic rhinitis: a systematic review and meta-analysis. *Am J Rhinol Allergy*. 2015 Jan-Feb;29(1):57-62.
55. Garcia MK, McQuade J, Lee R, Haddad R, Spano M, Cohen L. Acupuncture for symptom management in cancer care: an update. *Curr Oncol Rep*. 2014 Dec;16(12):418.
56. Cheong KB, Zhang JP, Huang Y, Zhang ZJ. The effectiveness of acupuncture in prevention and treatment of postoperative nausea and vomiting--a systematic review and meta-analysis. *PLOS ONE*. 2013;8(12):e82474.
57. Cho YH, Kim CK, Heo KH, Lee MS, Ha IH, Son DW, et al. Acupuncture for acute postoperative pain after back surgery: a systematic review and meta-analysis of randomized controlled trials. *Pain Pract*. 2015 Mar;15(3):279-91.
58. Barlow T, Downham C, Barlow D. The effect of complementary therapies on post-operative pain control in ambulatory knee surgery: a systematic review. *Complement Ther Med*. 2013 Oct;21(5):529-34.
59. Chen CC, Yang CC, Hu CC, Shih HN, Chang YH, Hsieh PH. Acupuncture for pain relief after total knee arthroplasty: a randomized controlled trial. *Reg Anesth Pain Med*. 2015 Jan-Feb;40(1):31-6.
60. Zhang R, Lao L, Ren K, Berman BM. Mechanisms of acupuncture-electroacupuncture on persistent pain. *Anesthesiology*. 2014 Feb;120(2):482-503.
61. Zhao ZQ. Neural mechanism underlying acupuncture analgesia. *Prog Neurobiol*. 2008 Aug;85(4):355-75.
62. Han JS. Acupuncture analgesia: areas of consensus and controversy. *Pain*. 2011 Mar;152(3 Suppl):S41-8.
63. Han JS. Acupuncture and endorphins. *Neurosci Lett*. 2004 May 6;361(1-3):258-61.
64. McDonald JL, Cripps AW, Smith PK. Mediators, Receptors, and Signalling Pathways in the Anti-Inflammatory and Antihyperalgesic Effects of Acupuncture. *Evid Based Complement Alternat Med*. 2015;2015:975632.
65. McDonald JL, Cripps AW, Smith PK, Smith CA, Xue CC, Golianu B. The anti-inflammatory effects of acupuncture and their relevance to allergic rhinitis: a narrative review and proposed model. *Evid Based Complement Alternat Med*. 2013;2013:591796.
66. Li YM, Zhuang LX, Lai XS, Jiang GH. [Effects of electroacupuncture on plasma vasoactive intestinal peptide and substance P in perennial allergic rhinitis patients]. *Zhen Ci Yan Jiu*. 2007 Apr;32(2):136-8.
67. Zhang Z, Wang C, Gu G, Li H, Zhao H, Wang K, et al. The effects of electroacupuncture at the ST36 (Zusanli) acupoint on cancer pain and transient receptor potential vanilloid subfamily 1 expression in Walker 256 tumor-bearing rats. *Anesth Analg*. 2012 Apr;114(4):879-85.
68. LU ZZ, Yin, X.J., Teng, W.J., Chen, Y.H., Sun, J., Zhao, J.M., Wang, A.Q. Bao C.H. & Shi, Y. Comparative effect of EA and moxibustion on the expression of SP and VIP in patients with IBS. *Journal of Traditional Chinese Medicine*. 2015;35(4):402-10.
69. Zhang YN, Zhao HJ, Wang Y, Lu Y, Wang SJ. [Effect of Electroacupuncture Intervention on Constipation-predominant Irritable Bowel Syndrome and Colonic CGRP and SP Expression in Rats]. *Zhen Ci Yan Jiu*. 2016 Feb;41(1):31-4.
70. Li W, Zhong, GW, Qi, M, Liu, WP, Wang, S, Wen, LB et al. Acupuncture treatment of migraine and plasma CGRP and SP content changes. *World Journal of Acupuncture-Moxibustion*. 2001;11(3):11-4.
71. Shi H, Li JH, Ji CF, Shang HY, Qiu EC, Wang JJ, et al. [Effect of electroacupuncture on cortical spreading depression and plasma CGRP and substance P contents in migraine rats]. *Zhen Ci Yan Jiu*. 2010 Feb;35(1):17-21.

72. Stener-Victorin E, Wu X. Effects and mechanisms of acupuncture in the reproductive system. *Auton Neurosci*. 2010 Oct 28;157(1-2):46-51.
73. Li H, He T, Xu Q, Li Z, Liu Y, Li F, et al. Acupuncture and regulation of gastrointestinal function. *World J Gastroenterol*. 2015 Jul 21;21(27):8304-13.
74. Qin Q, Mo Q, Liu K, He X, Gao X, Zhu B. Acupuncture at homotopic acupoints exerts dual effects on bladder motility in anesthetized rats. *BMC Complement Altern Med*. 2015;15:267.
75. Witt CM, Brinkhaus B. Efficacy, effectiveness and cost-effectiveness of acupuncture for allergic rhinitis - An overview about previous and ongoing studies. *Auton Neurosci*. 2010 Oct 28;157(1-2):42-5.
76. Kim SY, Lee H, Chae Y, Park HJ, Lee H. A systematic review of cost-effectiveness analyses alongside randomised controlled trials of acupuncture. *Acupunct Med*. 2012 Dec;30(4):273-85.
77. Reinhold T, Roll S, Willich SN, Ortiz M, Witt CM, Brinkhaus B. Cost-effectiveness for acupuncture in seasonal allergic rhinitis: economic results of the ACUSAR trial. *Ann Allergy Asthma Immunol*. 2013 Jul;111(1):56-63.
78. Reinhold T, Brinkhaus B, Willich SN, Witt C. Acupuncture in patients suffering from allergic asthma: is it worth additional costs? *J Altern Complement Med*. 2014 Mar;20(3):169-77.
79. Bazzan AJ, Zabrecky G, Monti DA, Newberg AB. Current evidence regarding the management of mood and anxiety disorders using complementary and alternative medicine. *Expert Rev Neurother*. 2014 Apr;14(4):411-23.
80. Goyata SL, Avelino CC, Santos SV, Souza Junior DI, Gurgel MD, Terra FS. Effects from acupuncture in treating anxiety: integrative review. *Rev Bras Enferm*. 2016 Jun;69(3):602-9.
81. Engel CC, Cordova EH, Benedek DM, Liu X, Gore KL, Goertz C, et al. Randomized effectiveness trial of a brief course of acupuncture for posttraumatic stress disorder. *Med Care*. 2014 Dec;52(12 Suppl 5):S57-64.
82. Bosch P, van den Noort M, Staudte H, Lim S. Schizophrenia and Depression: A systematic Review of the Effectiveness and the Working Mechanisms Behind Acupuncture. *Explore (NY)*. 2015 Jul-Aug;11(4):281-91.
83. Shen X, Xia J, Adams CE. Acupuncture for schizophrenia. *Cochrane Database Syst Rev*. 2014(10):Cd005475.
84. White AR, Rampes H, Liu JP, Stead LF, Campbell J. Acupuncture and related interventions for smoking cessation. *Cochrane Database Syst Rev*. 2014(1):Cd000009.
85. Grillo CM, Canales Gde L, Wada RS, Alves MC, Barbosa CM, Berzin F, et al. Could Acupuncture Be Useful in the Treatment of Temporomandibular Dysfunction? *J Acupunct Meridian Stud*. 2015 Aug;8(4):192-9.
86. Paley CA, Johnson MI, Tashani OA, Bagnall AM. Acupuncture for cancer pain in adults. *Cochrane Database Syst Rev*. 2015(10):Cd007753.
87. Chiu HY, Hsieh YJ, Tsai PS. Systematic review and meta-analysis of acupuncture to reduce cancer-related pain. *Eur J Cancer Care (Engl)*. 2016 Feb 7.
88. Chan YY, Lo WY, Yang SN, Chen YH, Lin JG. The benefit of combined acupuncture and antidepressant medication for depression: A systematic review and meta-analysis. *J Affect Disord*. 2015 May 1;176:106-17.
89. Spackman E, Richmond S, Sculpher M, Bland M, Brealey S, Gabe R, et al. Cost-effectiveness analysis of acupuncture, counselling and usual care in treating patients with depression: the results of the ACUDep trial. *PLOS ONE*. 2014;9(11):e113726.
90. Zhao K. Acupuncture for the treatment of insomnia. *Int Rev Neurobiol*. 2013;111:217-34.

91. Shergis JL, Ni X, Jackson ML, Zhang AL, Guo X, Li Y, et al. A systematic review of acupuncture for sleep quality in people with insomnia. *Complement Ther Med*. 2016 Jun;26:11-20.
92. Levett KM, Smith CA, Dahlen HG, Bensoussan A. Acupuncture and acupressure for pain management in labour and birth: a critical narrative review of current systematic review evidence. *Complement Ther Med*. 2014 Jun;22(3):523-40.
93. Selva Olid A, Martinez Zapata MJ, Sola I, Stojanovic Z, Uriona Tuma SM, Bonfill Cosp X. Efficacy and Safety of Needle Acupuncture for Treating Gynecologic and Obstetric Disorders: An Overview. *Med Acupunct*. 2013 Dec 1;25(6):386-97.
94. Close C, Sinclair M, Liddle SD, Madden E, McCullough JE, Hughes C. A systematic review investigating the effectiveness of Complementary and Alternative Medicine (CAM) for the management of low back and/or pelvic pain (LBPP) in pregnancy. *J Adv Nurs*. 2014 Aug;70(8):1702-16.
95. Liddle SD, Pennick V. Interventions for preventing and treating low-back and pelvic pain during pregnancy. *Cochrane Database Syst Rev*. 2015(9):Cd001139.
96. Chang SC, Hsu CH, Hsu CK, Yang SS, Chang SJ. The efficacy of acupuncture in managing patients with chronic prostatitis/chronic pelvic pain syndrome: A systemic review and meta-analysis. *Neurourol Urodyn*. 2016 Jan 6.
97. Qin Z, Wu J, Zhou J, Liu Z. Systematic Review of Acupuncture for Chronic Prostatitis/Chronic Pelvic Pain Syndrome. *Medicine (Baltimore)*. 2016 Mar;95(11):e3095.
98. Esteghamati A, Mazaheri T, Vahidi Rad M, Noshad S. Complementary and alternative medicine for the treatment of obesity: a critical review. *Int J Endocrinol Metab*. 2015 Apr;13(2):e19678.
99. Bega D, Malkani R. Alternative treatment of restless legs syndrome: an overview of the evidence for mind-body interventions, lifestyle interventions, and nutraceuticals. *Sleep Med*. 2016 Jan;17:99-105.
100. Smith CA, Crowther CA, Grant SJ. Acupuncture for induction of labour. *Cochrane Database Syst Rev*. 2013(8):Cd002962.
101. Mollart LJ, Adam J, Foureur M. Impact of acupressure on onset of labour and labour duration: A systematic review. *Women Birth*. 2015 Sep;28(3):199-206.
102. Li P, Qiu T, Qin C. Efficacy of Acupuncture for Bell's Palsy: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *PLOS ONE*. 2015;10(5):e0121880.
103. Kwon HJ, Choi JY, Lee MS, Kim YS, Shin BC, Kim JI. Acupuncture for the sequelae of Bell's palsy: a randomized controlled trial. *Trials*. 2015;16:246.
104. Law SK, Li T. Acupuncture for glaucoma. *Cochrane Database Syst Rev*. 2013(5):Cd006030.
105. Dalton-Brewer N. The Role of Complementary and Alternative Medicine for the Management of Fibroids and Associated Symptomatology. *Curr Obstet Gynecol Rep*. 2016;5:110-8.
106. Cao H, Wang Y, Chang D, Zhou L, Liu J. Acupuncture for vascular mild cognitive impairment: a systematic review of randomised controlled trials. *Acupunct Med*. 2013 Dec;31(4):368-74.
107. Cui X, Zhou J, Qin Z, Liu Z. Acupuncture for Erectile Dysfunction: A Systematic Review. *Biomed Res Int*. 2016;2016:2171923.
108. He M, Li X, Liu Y, Zhong J, Jiang L, Liu Y, et al. Electroacupuncture for Tinnitus: A Systematic Review. *PLOS ONE*. 2016;11(3):e0150600.

109. Wu SL, Leung AW, Yew DT. Acupuncture for Detoxification in Treatment of Opioid Addiction. *East Asian Arch Psychiatry*. 2016 Jun;26(2):70-6.
110. Grant S, Kandrack R, Motala A, Shanman R, Booth M, Miles J, et al. Acupuncture for substance use disorders: A systematic review and meta-analysis. *Drug Alcohol Depend*. 2016 Jun 1;163:1-15.
111. Kim JE, Seo BK, Choi JB, Kim HJ, Kim TH, Lee MH, et al. Acupuncture for chronic fatigue syndrome and idiopathic chronic fatigue: a multicenter, nonblinded, randomized controlled trial. *Trials*. 2015;16:314.
112. Deare JC, Zheng Z, Xue CC, Liu JP, Shang J, Scott SW, et al. Acupuncture for treating fibromyalgia. *Cochrane Database Syst Rev*. 2013(5):Cd007070.
113. Yang B, Yi G, Hong W, Bo C, Wang Z, Liu Y, et al. Efficacy of acupuncture on fibromyalgia syndrome: a meta-analysis. *J Tradit Chin Med*. 2014 Aug;34(4):381-91.
114. Lauche R, Cramer H, Hauser W, Dobos G, Langhorst J. A Systematic Overview of Reviews for Complementary and Alternative Therapies in the Treatment of the Fibromyalgia Syndrome. *Evid Based Complement Alternat Med*. 2015;2015:610615.
115. Jang SH, Kim DI, Choi MS. Effects and treatment methods of acupuncture and herbal medicine for premenstrual syndrome/premenstrual dysphoric disorder: systematic review. *BMC Complement Altern Med*. 2014;14:11.
116. Hofmeister S, Bodden S. Premenstrual Syndrome and Premenstrual Dysphoric Disorder. *Am Fam Physician*. 2016 Aug 1;94(3):236-40.
117. Cheong YC, Dix S, Hung Yu Ng E, Ledger WL, Farquhar C. Acupuncture and assisted reproductive technology. *Cochrane Database Syst Rev*. 2013(7):Cd006920.
118. Shen C, Wu M, Shu D, Zhao X, Gao Y. The role of acupuncture in in vitro fertilization: a systematic review and meta-analysis. *Gynecol Obstet Invest*. 2015;79(1):1-12.
119. Qian Y, Xia XR, Ochin H, Huang C, Gao C, Gao L, et al. Therapeutic effect of acupuncture on the outcomes of in vitro fertilization: a systematic review and meta-analysis. *Arch Gynecol Obstet*. 2016 Dec 19.
120. Jo J, Lee YJ. Effectiveness of acupuncture in women with polycystic ovarian syndrome undergoing in vitro fertilisation or intracytoplasmic sperm injection: a systematic review and meta-analysis. *Acupunct Med*. 2017 Jan 11.
121. Fernandez-Llanio Comella N, Fernandez Matilla M, Castellano Cuesta JA. Have complementary therapies demonstrated effectiveness in rheumatoid arthritis? *Reumatol Clin*. 2016 May-Jun;12(3):151-7.
122. Matthews A, Haas DM, O'Mathuna DP, Dowswell T. Interventions for nausea and vomiting in early pregnancy. *Cochrane Database Syst Rev*. 2015(9):Cd007575.
123. Cheuk DK, Wong V. Acupuncture for epilepsy. *Cochrane Database Syst Rev*. 2014(5):Cd005062.
124. Lardon A, Girard MP, Zaim C, Lemeunier N, Descarreaux M, Marchand AA. Effectiveness of preventive and treatment interventions for primary headaches in the workplace: A systematic review of the literature. *Cephalalgia*. 2016 Mar 2.
125. Cho HK, Park IJ, Jeong YM, Lee YJ, Hwang SH. Can perioperative acupuncture reduce the pain and vomiting experienced after tonsillectomy? A meta-analysis. *Laryngoscope*. 2016 Mar;126(3):608-15.
126. Shin HC, Kim JS, Lee SK, Kwon SH, Kim MS, Lee EJ, et al. The effect of acupuncture on postoperative nausea and vomiting after pediatric tonsillectomy: A meta-analysis and systematic review. *Laryngoscope*. 2016 Aug;126(8):1761-7.

127. Zhang JH, Wang D, Liu M. Overview of systematic reviews and meta-analyses of acupuncture for stroke. *Neuroepidemiology*. 2014;42(1):50-8.
128. Vados L, Ferreira A, Zhao S, Vercelino R, Wang S. Effectiveness of acupuncture combined with rehabilitation for treatment of acute or subacute stroke: a systematic review. *Acupunct Med*. 2015 Jun;33(3):180-7.
129. Liu AJ, Li JH, Li HQ, Fu DL, Lu L, Bian ZX, et al. Electroacupuncture for Acute Ischemic Stroke: A Meta-Analysis of Randomized Controlled Trials. *Am J Chin Med*. 2015;43(8):1541-66.
130. Yang A, Wu HM, Tang JL, Xu L, Yang M, Liu GJ. Acupuncture for stroke rehabilitation. *Cochrane Database Syst Rev*. 2016 Aug 26;8:Cd004131.
131. Lim SM, Yoo J, Lee E, Kim HJ, Shin S, Han G, et al. Acupuncture for spasticity after stroke: a systematic review and meta-analysis of randomized controlled trials. *Evid Based Complement Alternat Med*. 2015;2015:870398.
132. Rodriguez-Mansilla J, Espejo-Antunez L, Bustamante-Lopez AI. [Effectiveness of acupuncture in spasticity of the post-stroke patient. Systematic review]. *Aten Primaria*. 2016 Apr;48(4):226-34.
133. Lee SH, Lim SM. Acupuncture for insomnia after stroke: a systematic review and meta-analysis. *BMC Complement Altern Med*. 2016;16:228.
134. Lee SH, Lim SM. Acupuncture for Poststroke Shoulder Pain: A Systematic Review and Meta-Analysis. *Evid Based Complement Alternat Med*. 2016;2016:3549878.
135. Yue J, Liu M, Li J, Wang Y, Hung ES, Tong X, et al. Acupuncture for the treatment of hiccups following stroke: a systematic review and meta-analysis. *Acupunct Med*. 2016 Jun 10.
136. Chiu HY, Pan CH, Shyu YK, Han BC, Tsai PS. Effects of acupuncture on menopause-related symptoms and quality of life in women in natural menopause: a meta-analysis of randomized controlled trials. *Menopause*. 2015 Feb;22(2):234-44.
137. Dodin S, Blanchet C, Marc I, Ernst E, Wu T, Vaillancourt C, et al. Acupuncture for menopausal hot flashes. *Cochrane Database Syst Rev*. 2013(7):Cd007410.
138. Chiu HY, Hsieh YJ, Tsai PS. Acupuncture to Reduce Sleep Disturbances in Perimenopausal and Postmenopausal Women: A Systematic Review and Meta-analysis. *Obstet Gynecol*. 2016 Mar;127(3):507-15.
139. Bezerra AG, Pires GN, Andersen ML, Tufik S, Hachul H. Acupuncture to Treat Sleep Disorders in Postmenopausal Women: A Systematic Review. *Evid Based Complement Alternat Med*. 2015;2015:563236.
140. Lau CH, Wu X, Chung VC, Liu X, Hui EP, Cramer H, et al. Acupuncture and Related Therapies for Symptom Management in Palliative Cancer Care: Systematic Review and Meta-Analysis. *Medicine (Baltimore)*. 2016 Mar;95(9):e2901.
141. Lu W, Rosenthal DS. Acupuncture for cancer pain and related symptoms. *Curr Pain Headache Rep*. 2013 Mar;17(3):321.
142. Towler P, Molassiotis A, Brearley SG. What is the evidence for the use of acupuncture as an intervention for symptom management in cancer supportive and palliative care: an integrative overview of reviews. *Support Care Cancer*. 2013 Oct;21(10):2913-23.
143. Lian WL, Pan MQ, Zhou DH, Zhang ZJ. Effectiveness of acupuncture for palliative care in cancer patients: a systematic review. *Chin J Integr Med*. 2014 Feb;20(2):136-47.
144. He XR, Wang Q, Li PP. Acupuncture and moxibustion for cancer-related fatigue: a systematic review and meta-analysis. *Asian Pac J Cancer Prev*. 2013;14(5):3067-74.

145. Ling WM, Lui LY, So WK, Chan K. Effects of acupuncture and acupressure on cancer-related fatigue: a systematic review. *Oncol Nurs Forum*. 2014 Nov 1;41(6):581-92.
146. Posadzki P, Moon TW, Choi TY, Park TY, Lee MS, Ernst E. Acupuncture for cancer-related fatigue: a systematic review of randomized clinical trials. *Support Care Cancer*. 2013 Jul;21(7):2067-73.
147. Finnegan-John J, Molassiotis A, Richardson A, Ream E. A systematic review of complementary and alternative medicine interventions for the management of cancer-related fatigue. *Integr Cancer Ther*. 2013 Jul;12(4):276-90.
148. Zeng Y, Luo T, Finnegan-John J, Cheng AS. Meta-Analysis of Randomized Controlled Trials of Acupuncture for Cancer-Related Fatigue. *Integr Cancer Ther*. 2014 May;13(3):193-200.
149. Choi TY, Kim JI, Lim HJ, Lee MS. Acupuncture for Managing Cancer-Related Insomnia: A Systematic Review of Randomized Clinical Trials. *Integr Cancer Ther*. 2016 Aug 16.
150. Garcia MK, Graham-Getty L, Haddad R, Li Y, McQuade J, Lee RT, et al. Systematic review of acupuncture to control hot flashes in cancer patients. *Cancer*. 2015 Nov 15;121(22):3948-58.
151. Johns C, Seav SM, Dominick SA, Gorman JR, Li H, Natarajan L, et al. Informing hot flash treatment decisions for breast cancer survivors: a systematic review of randomized trials comparing active interventions. *Breast Cancer Res Treat*. 2016 Apr;156(3):415-26.
152. Chen YP, Liu T, Peng YY, Wang YP, Chen H, Fan YF, et al. Acupuncture for hot flashes in women with breast cancer: A systematic review. *J Cancer Res Ther*. 2016 Apr-Jun;12(2):535-42.
153. Salehi A, Marzban M, Zadeh AR. Acupuncture for treating hot flashes in breast cancer patients: an updated meta-analysis. *Support Care Cancer*. 2016 Aug 6.
154. Zhuang L, Yang Z, Zeng X, Zhua X, Chen Z, Liu L, et al. The preventive and therapeutic effect of acupuncture for radiation-induced xerostomia in patients with head and neck cancer: a systematic review. *Integr Cancer Ther*. 2013 May;12(3):197-205.
155. Furness S, Bryan G, McMillan R, Worthington HV. Interventions for the management of dry mouth: non-pharmacological interventions. *Cochrane Database Syst Rev*. 2013(8):Cd009603.
156. Hanchanale S, Adkinson L, Daniel S, Fleming M, Oxberry SG. Systematic literature review: xerostomia in advanced cancer patients. *Support Care Cancer*. 2015 Mar;23(3):881-8.
157. Kim KH, Kim DH, Kim HY, Son GM. Acupuncture for recovery after surgery in patients undergoing colorectal cancer resection: a systematic review and meta-analysis. *Acupunct Med*. 2016 Aug;34(4):248-56.
158. Bae K, Yoo HS, Lamoury G, Boyle F, Rosenthal DS, Oh B. Acupuncture for Aromatase Inhibitor-Induced Arthralgia: A Systematic Review. *Integr Cancer Ther*. 2015 Nov;14(6):496-502.
159. Chien TJ, Liu CY, Chang YF, Fang CJ, Hsu CH. Acupuncture for treating aromatase inhibitor-related arthralgia in breast cancer: a systematic review and meta-analysis. *J Altern Complement Med*. 2015 May;21(5):251-60.
160. Franconi G, Manni L, Schroder S, Marchetti P, Robinson N. A systematic review of experimental and clinical acupuncture in chemotherapy-induced peripheral neuropathy. *Evid Based Complement Alternat Med*. 2013;2013:516916.
161. Fu H, Chen B, Hong S, Guo Y. Acupuncture Therapy for the Treatment of Myelosuppression after Chemotherapy: A Literature Review over the Past 10 Years. *J Acupunct Meridian Stud*. 2015 Jun;8(3):122-6.
162. Haddad NE, Palesh O. Acupuncture in the treatment of cancer-related psychological symptoms. *Integr Cancer Ther*. 2014 Sep;13(5):371-85.

163. Lee MS, Ernst E. Acupuncture for surgical conditions: an overview of systematic reviews. *Int J Clin Pract.* 2014 Jun;68(6):783-9.
164. Asmussen S, Maybauer DM, Chen JD, Fraser JF, Toon MH, Przkora R, et al. Effects of Acupuncture in Anesthesia for Craniotomy: A Meta-Analysis. *J Neurosurg Anesthesiol.* 2016 Mar 10.
165. Liodden I, Norheim AJ. Acupuncture and related techniques in ambulatory anesthesia. *Curr Opin Anaesthesiol.* 2013 Dec;26(6):661-8.
166. Yang L, Yang Z, Yu H, Song H. Acupuncture therapy is more effective than artificial tears for dry eye syndrome: evidence based on a meta-analysis. *Evid Based Complement Alternat Med.* 2015;2015:143858.
167. Hadianfard M, Bazrafshan E, Momeninejad H, Jahani N. Efficacies of Acupuncture and Anti-inflammatory Treatment for Carpal Tunnel Syndrome. *J Acupunct Meridian Stud.* 2015 Oct;8(5):229-35.
168. Manheimer E, Wieland LS, Cheng K, Li SM, Shen X, Berman BM, et al. Acupuncture for irritable bowel syndrome: systematic review and meta-analysis. *Am J Gastroenterol.* 2012 Jun;107(6):835-47; quiz 48.
169. MacPherson H, Tilbrook H, Agbedjro D, Buckley H, Hewitt C, Frost C. Acupuncture for irritable bowel syndrome: 2-year follow-up of a randomised controlled trial. *Acupunct Med.* 2016 Mar 15.
170. Wang J, Xiong X, Liu W. Acupuncture for essential hypertension. *Int J Cardiol.* 2013 Nov 20;169(5):317-26.
171. Li DZ, Zhou Y, Yang YN, Ma YT, Li XM, Yu J, et al. Acupuncture for essential hypertension: a meta-analysis of randomized sham-controlled clinical trials. *Evid Based Complement Alternat Med.* 2014;2014:279478.
172. Zhao XF, Hu HT, Li JS, Shang HC, Zheng HZ, Niu JF, et al. Is Acupuncture Effective for Hypertension? A Systematic Review and Meta-Analysis. *PLOS ONE.* 2015;10(7):e0127019.
173. Gadau M, Yeung WF, Liu H, Zaslowski C, Tan YS, Wang FC, et al. Acupuncture and moxibustion for lateral elbow pain: a systematic review of randomized controlled trials. *BMC Complement Altern Med.* 2014;14:136.
174. Tang H, Fan H, Chen J, Yang M, Yi X, Dai G, et al. Acupuncture for Lateral Epicondylitis: A Systematic Review. *Evid Based Complement Alternat Med.* 2015;2015:861849.
175. van der Velde G, Yu H, Paulden M, Cote P, Varatharajan S, Shearer HM, et al. Which interventions are cost-effective for the management of whiplash-associated and neck pain-associated disorders? A systematic review of the health economic literature by the Ontario Protocol for Traffic Injury Management (OPTIMa) Collaboration. *Spine J.* 2015 Nov 26.
176. Trinh K, Graham N, Irnich D, Cameron ID, Forget M. Acupuncture for neck disorders. *Cochrane Database Syst Rev.* 2016(5):Cd004870.
177. Moon TW, Posadzki P, Choi TY, Park TY, Kim HJ, Lee MS, et al. Acupuncture for treating whiplash associated disorder: a systematic review of randomised clinical trials. *Evid Based Complement Alternat Med.* 2014;2014:870271.
178. Dong W, Goost H, Lin XB, Burger C, Paul C, Wang ZL, et al. Treatments for shoulder impingement syndrome: a PRISMA systematic review and network meta-analysis. *Medicine (Baltimore).* 2015 Mar;94(10):e510.
179. Liu CF, Chien LW. Efficacy of acupuncture in children with asthma: a systematic review. *Ital J Pediatr.* 2015;41:48.
180. Lee SH, Chang GT, Zhang X, Lee H. Acupoint Herbal Patching for Asthma: A Systematic Review and Meta-analysis of Randomized Controlled Trials. *Medicine (Baltimore).* 2016 Jan;95(2):e2439.

181. Su L, Meng L, Chen R, Wu W, Peng B, Man L. Acupoint Application for Asthma Therapy in Adults: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Forsch Komplementmed*. 2016;23(1):16-21.
182. Chen MN, Chien LW, Liu CF. Acupuncture or Acupressure at the Sanyinjiao (SP6) Acupoint for the Treatment of Primary Dysmenorrhea: A Meta-Analysis. *Evid Based Complement Alternat Med*. 2013;2013:493038.
183. Xu T, Hui L, Juan YL, Min SG, Hua WT. Effects of moxibustion or acupoint therapy for the treatment of primary dysmenorrhea: a meta-analysis. *Altern Ther Health Med*. 2014 Jul-Aug;20(4):33-42.
184. Abaraogu UO, Tabansi-Ochuogu CS. As Acupressure Decreases Pain, Acupuncture May Improve Some Aspects of Quality of Life for Women with Primary Dysmenorrhea: A Systematic Review with Meta-Analysis. *J Acupunct Meridian Stud*. 2015 Oct;8(5):220-8.
185. Smith CA, Armour M, Zhu X, Li X, Lu ZY, Song J. Acupuncture for dysmenorrhoea. *Cochrane Database Syst Rev*. 2016;4:Cd007854.
186. Kim TH, Lee MS, Kim KH, Kang JW, Choi TY, Ernst E. Acupuncture for treating acute ankle sprains in adults. *Cochrane Database Syst Rev*. 2014(6):Cd009065.
187. Xu L, Xu H, Gao W, Wang W, Zhang H, Lu DP. Treating angina pectoris by acupuncture therapy. *Acupunct Electrother Res*. 2013;38(1-2):17-35.
188. Yu C, Ji K, Cao H, Wang Y, Jin HH, Zhang Z, et al. Effectiveness of acupuncture for angina pectoris: a systematic review of randomized controlled trials. *BMC Complement Altern Med*. 2015;15:90.
189. Zhang Z, Bai R, Zhang L, Qi W, Wang Y, Li B, et al. [Acupuncture combined with Western medicine for angina of coronary artery disease: a systematic review]. *Zhongguo Zhen Jiu*. 2015 Apr;35(4):407-11.
190. Coyle ME, Shergis JL, Huang ET, Guo X, Di YM, Zhang A, et al. Acupuncture therapies for chronic obstructive pulmonary disease: a systematic review of randomized, controlled trials. *Altern Ther Health Med*. 2014 Nov-Dec;20(6):10-23.
191. Lee H, Kim TH, Leem J. Acupuncture for heart failure: A systematic review of clinical studies. *Int J Cardiol*. 2016 Jul 30;222:321-31.
192. Min D, Xu-Feng W. An Updated Meta-Analysis of the Efficacy and Safety of Acupuncture Treatment for Vascular Cognitive Impairment Without Dementia. *Curr Neurovasc Res*. 2016;13(3):230-8.
193. Park J, Hahn S, Park JY, Park HJ, Lee H. Acupuncture for ankle sprain: systematic review and meta-analysis. *BMC Complement Altern Med*. 2013;13:55.
194. Kim KH, Kim TH, Lee BR, Kim JK, Son DW, Lee SW, et al. Acupuncture for lumbar spinal stenosis: a systematic review and meta-analysis. *Complement Ther Med*. 2013 Oct;21(5):535-56.
195. Nahin RL, Boineau R, Khalsa PS, Stussman BJ, Weber WJ. Evidence-Based Evaluation of Complementary Health Approaches for Pain Management in the United States. *Mayo Clin Proc*. 2016 Sep;91(9):1292-306.
196. Lewis RA, Williams NH, Sutton AJ, Burton K, Din NU, Matar HE, et al. Comparative clinical effectiveness of management strategies for sciatica: systematic review and network meta-analyses. *Spine J*. 2015 Jun 1;15(6):1461-77.
197. Qin Z, Liu X, Wu J, Zhai Y, Liu Z. Effectiveness of Acupuncture for Treating Sciatica: A Systematic Review and Meta-Analysis. *Evid Based Complement Alternat Med*. 2015;2015:425108.
198. Ji M, Wang X, Chen M, Shen Y, Zhang X, Yang J. The Efficacy of Acupuncture for the Treatment of Sciatica: A Systematic Review and Meta-Analysis. *Evid Based Complement Alternat Med*. 2015;2015:192808.

199. Kim HJ, Jeon BS. Is acupuncture efficacious therapy in Parkinson's disease? *J Neurol Sci.* 2014 Jun 15;341(1-2):1-7.
200. Zhou J, Peng W, Xu M, Li W, Liu Z. The effectiveness and safety of acupuncture for patients with Alzheimer disease: a systematic review and meta-analysis of randomized controlled trials. *Medicine (Baltimore).* 2015 Jun;94(22):e933.
201. Au DW, Tsang HW, Ling PP, Leung CH, Ip PK, Cheung WM. Effects of acupressure on anxiety: a systematic review and meta-analysis. *Acupunct Med.* 2015 Oct;33(5):353-9.
202. Chen HM, Chen CH. Effects of acupressure at the Sanyinjiao point on primary dysmenorrhoea. *J Adv Nurs.* 2004 Nov;48(4):380-7.
203. Boelig RC, Barton SJ, Saccone G, Kelly AJ, Edwards SJ, Berghella V. Interventions for treating hyperemesis gravidarum. *Cochrane Database Syst Rev.* 2016(5):Cd010607.
204. Chai Q, Fei Y, Cao H, Wang C, Tian J, Liu J. Acupuncture for melasma in women: a systematic review of randomised controlled trials. *Acupunct Med.* 2015 Aug;33(4):254-61.
205. Manheimer E, van der Windt D, Cheng K, Stafford K, Liu J, Tierney J, et al. The effects of acupuncture on rates of clinical pregnancy among women undergoing in vitro fertilization: a systematic review and meta-analysis. *Hum Reprod Update.* 2013 Nov-Dec;19(6):696-713.
206. Kwan I, Bhattacharya S, Knox F, McNeil A. Pain relief for women undergoing oocyte retrieval for assisted reproduction. *Cochrane Database Syst Rev.* 2013(1):Cd004829.
207. Ren LN, Guo LH, Ma WZ, Zhang R. [A meta-analysis on acupuncture treatment of polycystic ovary syndrome]. *Zhen Ci Yan Jiu.* 2014 Jun;39(3):238-46.
208. Lim CE, Ng RW, Xu K, Cheng NC, Xue CC, Liu JP, et al. Acupuncture for polycystic ovarian syndrome. *Cochrane Database Syst Rev.* 2016(5):Cd007689.
209. Jo J, Lee YJ, Lee H. Effectiveness of Acupuncture for Primary Ovarian Insufficiency: A Systematic Review and Meta-Analysis. *Evid Based Complement Alternat Med.* 2015;2015:842180.
210. Yang M, Li X, Liu S, Li Z, Xue M, Gao D, et al. Meta-analysis of acupuncture for relieving non-organic dyspeptic symptoms suggestive of diabetic gastroparesis. *BMC Complement Altern Med.* 2013;13:311.
211. Lan L, Zeng F, Liu GJ, Ying L, Wu X, Liu M, et al. Acupuncture for functional dyspepsia. *Cochrane Database Syst Rev.* 2014(10):Cd008487.
212. Kim KN, Chung SY, Cho SH. Efficacy of acupuncture treatment for functional dyspepsia: A systematic review and meta-analysis. *Complement Ther Med.* 2015 Dec;23(6):759-66.
213. Langhorst J, Wulfert H, Lauche R, Klose P, Cramer H, Dobos GJ, et al. Systematic review of complementary and alternative medicine treatments in inflammatory bowel diseases. *J Crohns Colitis.* 2015 Jan;9(1):86-106.
214. Kim KH, Lee MS, Kim TH, Kang JW, Choi TY, Lee JD. Acupuncture and related interventions for symptoms of chronic kidney disease. *Cochrane Database Syst Rev.* 2016(6):Cd009440.
215. Jerng UM, Jo JY, Lee S, Lee JM, Kwon O. The effectiveness and safety of acupuncture for poor semen quality in infertile males: a systematic review and meta-analysis. *Asian J Androl.* 2014 Nov-Dec;16(6):884-91.
216. Wang Y, Zhishun L, Peng W, Zhao J, Liu B. Acupuncture for stress urinary incontinence in adults. *Cochrane Database Syst Rev.* 2013(7):Cd009408.

217. Paik SH, Han SR, Kwon OJ, Ahn YM, Lee BC, Ahn SY. Acupuncture for the treatment of urinary incontinence: A review of randomized controlled trials. *Exp Ther Med*. 2013 Sep;6(3):773-80.
218. Naik PN, Kiran RA, Yalamanchal S, Kumar VA, Goli S, Vashist N. Acupuncture: An Alternative Therapy in Dentistry and Its Possible Applications. *Med Acupunct*. 2014 Dec 1;26(6):308-14.
219. Cheong KB, Zhang JP, Huang Y. The effectiveness of acupuncture in postoperative gastroparesis syndrome--a systematic review and meta-analysis. *Complement Ther Med*. 2014 Aug;22(4):767-86.
220. Cheong KB, Zhang J, Huang Y. [Effectiveness of acupuncture in postoperative ileus: a systematic review and Meta-analysis]. *J Tradit Chin Med*. 2016 Jun;36(3):271-82.
221. Hu C, Zhang H, Wu W, Yu W, Li Y, Bai J, et al. Acupuncture for Pain Management in Cancer: A Systematic Review and Meta-Analysis. *Evid Based Complement Alternat Med*. 2016;2016:1720239.
222. Xue CC, Zhang AL, Zhang CS, DaCosta C, Story DF, Thien FC. Acupuncture for seasonal allergic rhinitis: a randomized controlled trial. *Ann Allergy Asthma Immunol*. 2015 Oct;115(4):317-24.e1.
223. He J, Jiang L, Peng T, Xia M, Chen H. Acupuncture Points Stimulation for Meniere's Disease/Syndrome: A Promising Therapeutic Approach. *Evid Based Complement Alternat Med*. 2016;2016:6404197.
224. Zhang XC, Xu XP, Xu WT, Hou WZ, Cheng YY, Li CX, et al. Acupuncture therapy for sudden sensorineural hearing loss: a systematic review and meta-analysis of randomized controlled trials. *PLOS ONE*. 2015;10(4):e0125240.
225. Li S, Yu B, Zhou D, He C, Kang L, Wang X, et al. Acupuncture for Attention Deficit Hyperactivity Disorder (ADHD) in children and adolescents. *Cochrane Database Syst Rev*. 2011(4):Cd007839.
226. Cheuk DK, Wong V, Chen WX. Acupuncture for autism spectrum disorders (ASD). *Cochrane Database Syst Rev*. 2011(9):Cd007849.
227. Wong V, Cheuk DK, Chu V. Acupuncture for hypoxic ischemic encephalopathy in neonates. *Cochrane Database Syst Rev*. 2013(1):Cd007968.
228. He J, Jia P, Zheng M, Zhang M, Jiang H. Acupuncture for mumps in children. *Cochrane Database Syst Rev*. 2015(2):Cd008400.
229. Ma C, Sivamani RK. Acupuncture as a Treatment Modality in Dermatology: A Systematic Review. *J Altern Complement Med*. 2015 Sep;21(9):520-9.
230. Tan HY, Lenon GB, Zhang AL, Xue CC. Efficacy of acupuncture in the management of atopic dermatitis: a systematic review. *Clin Exp Dermatol*. 2015 Oct;40(7):711-5; quiz 5-6.
231. Vieira BL, Lim NR, Lohman ME, Lio PA. Complementary and Alternative Medicine for Atopic Dermatitis: An Evidence-Based Review. *Am J Clin Dermatol*. 2016 Jul 7.
232. Yao Q, Li S, Liu X, Qin Z, Liu Z. The Effectiveness and Safety of Acupuncture for Patients with Chronic Urticaria: A Systematic Review. *Biomed Res Int*. 2016;2016:5191729.
233. Yu C, Zhang P, Lv ZT, Li JJ, Li HP, Wu CH, et al. Efficacy of Acupuncture in Itch: A Systematic Review and Meta-Analysis of Clinical Randomized Controlled Trials. *Evid Based Complement Alternat Med*. 2015;2015:208690.
234. Coyle M, Deng J, Zhang AL, Yu J, Guo X, Xue CC, et al. Acupuncture therapies for psoriasis vulgaris: a systematic review of randomized controlled trials. *Forsch Komplementmed*. 2015;22(2):102-9.
235. Kim KH, Lee BR, Ryu JH, Choi TY, Yang GY. The role of acupuncture in emergency department settings: a systematic review. *Complement Ther Med*. 2013 Feb;21(1):65-72.

236. del Pino-Sedeno T, Trujillo-Martin MM, Ruiz-Irastorza G, Cuellar-Pompa L, de Pascual-Medina AM, Serrano-Aguilar P. Effectiveness of Nonpharmacologic Interventions for Decreasing Fatigue in Adults With Systemic Lupus Erythematosus: A Systematic Review. *Arthritis Care Res (Hoboken)*. 2016 Jan;68(1):141-8.
237. Hackett KL, Deane KH, Strassheim V, Deary V, Rapley T, Newton JL, et al. A systematic review of non-pharmacological interventions for primary Sjogren's syndrome. *Rheumatology (Oxford)*. 2015 Nov;54(11):2025-32.
238. Baeumler PI, Fleckenstein J, Takayama S, Simang M, Seki T, Irnich D. Effects of acupuncture on sensory perception: a systematic review and meta-analysis. *PLOS ONE*. 2014;9(12):e113731.
239. MacPherson H, Vertosick EA, Foster NE, Lewith G, Linde K, Sherman KJ, et al. The persistence of the effects of acupuncture after a course of treatment: A meta-analysis of patients with chronic pain. *Pain*. 2016 Oct 17.
240. Zhang J, Shang H, Gao X, Ernst E. Acupuncture-related adverse events: a systematic review of the Chinese literature. *Bulletin of the World Health Organization*. [Review]. 2010 Dec 1;88(12):915-21C.
241. National Centre for Complementary and Alternative Health. News Releases - NIH review finds nondrug approaches effective for treatment of common pain conditions. US Department of Health and Human Services; 2016 [cited 2016]; Available from: <https://www.nih.gov/news-events/news-releases/nihreview-finds-nondrug-approaches-effective-treatment-common-pain-conditions>.
242. National Health and Medical Research Council. NHMRC additional levels of evidence and grades for recommendations for developers of guidelines. Table 3. NHMRC Evidence Hierarchy: designations of 'levels of evidence' according to type of research question (including explanatory notes). National Health and Medical Research Council; 2009.
243. Balshem H, Helfand M, Schunemann HJ, Oxman AD, Kunz R, Brozek J, et al. GRADE guidelines: 3. Rating the quality of evidence. Table 2. Significance of the four levels of evidence. *J Clin Epidemiol*. 2011 Apr;64(4):404.